Population characteristics of the spiny dogfish, *Squalus acanthias* Linnaeus, 1758, from geographically distinct locations in Atlantic Canada during the summer and fall of 1996

by

Tina Mae Moore

B. Sc. (Hon.), Acadia University, 1995

Thesis submitted in partial fulfillment of the requirements for the Degree of Master of Science (Biology)

> Acadia University Fall Convocation 1998

© by Tina Mae Moore, 1998

I, Tina Mae Moore, grant permission to the University Librarian at Acadia University to reproduce, loan, or distrubute copies of my thesis in microform, paper or electronic formats on a non-profit basis. I, however, retain the copyright in my thesis.

Signature of Author

Date

#### **Table of Contents**

Page	e
Introduction	
Materials and Methods	
Location	
Capture Methods and Measurements	
Mark and Recapture Methods	
Spine Removals and Ageing Techniques.	
Calculations of Worn Spine Age and Growth.	
Reproductive Tract Removal and Measurements	
Fecundity, Maturity, and Embryo Development	
Analysis of Variance of Total Length, Age, and Maturity	
of Dogfish in the Three Study Locations	
Estimation of Growth Parameters	
Results	
General Population Characteristics	
Weight-Length Relationship and Growth Comparisons	
Age and Growth	
Maturity	
Fecundity and Reproductive Development	
Migration	

Discussion	28
General Population Characteristics.	28
Growth	29
	31
Reproduction and Maturity	33
Management of Spiny Dogfish.	35
Bibliography	40
Tables	51
Figures	60
Appendices	23

#### Lists of Tables

3

#### Page

Table 6: Percent number of female dogfish in the reproductive	
development stages from the three study locations	
during the summer - fall 1996	57

Table 7: Summary of egg diameter range (mm), embryo total
length range (mm), and male and female dogfish pups
total length range (mm) in the various reproductive
development stages

# List of Figures

Page	9
Figure 1: The Minas Basin sampling locations for	
dogfish captured from July 7, 1996 to October 10,	
1996	
Figure 2: The N246 DFO survey area and set locations for spiny	
dogfish captured during July 1996	
Figure 3: The N249 DFO survey area and set locations for dogfish	
captured during September 1996	
Figure 4: Posterior dorsal spine removal in the spiny dogfish for	
an age estimate. a = first cut; b = second cut	
Figure 5: Longitudinal and schematic diagram of the posterior	
dorsal spine (a) and spine growth showing five annual	
opaque rings or bands at the cone bases (b)	
(Slauson 1982)	

Figure 6: Measurements taken from the posterior dorsal spine		
of the spiny dogfish. SL = Spine Length; WL = Wear		
Length; SBD = Spine Base Diameter; and WPD =		
Wear Point Diameter		
Figure 7: Reproductive tract removal in the female spiny		
dogfish		
Figure 8: Stage I of reproductive tract development in the		
female spiny dogfish		
Figure 9: Stage II of reproductive tract development in the		
female spiny dogfish		
Figure 10: Stage III of reproductive tract development in the		
female spiny dogfish		
Figure 11: Stage IV of reproductive tract development in the		
female spiny dogfish		

# Figure 12: Stage V of reproductive tract development in the Figure 13: Stage VI of reproductive tract development in the Figure 14: The N246 DFO Scotian Shelf survey set locations and number of spiny dogfish captured in each tow during Figure 15: The N249 DFO Gulf of St. Lawrence survey set locations and number of spiny dogfish captured in each Figure 16: Total length frequency of female dogfish captured in the Minas Basin, during the N246 DFO survey, and during the N249 DFO survey, summer - fall 1996. . . . . . . . 92

Figure 18: Age frequency of female dogfish captured in the Minas		
Basin, during the N246 DFO survey, and during the N249		
DFO survey, summer - fall 1996	6	

Figure 20:	Weight-length relationship of female spiny dogfish
	captured in the Minas Basin, during the N246 DFO
	survey, and during the N249 DFO survey,
	summer - fall 1996

Figure 23:	Spine base diameter (mm) in relation to age (years) of
	male and female dogfish with wear point diameters
	$\leq$ 1.5 mm to determine the equation for lost annuli due
	to wear

Figure 24:	The von Bertalanffy growth equations for female	
	dogfish captured in the Minas Basin, during the N246	
	DFO survey, and during the N249 DFO survey,	
	summer - fall 1996. Bars represent 1 standard deviation	
	from the mean	108

# 

Figure 27: Percent maturity of female	e dogfish captured in the
Minas Basin, during the N	1246 DFO survey, and during
the N249 DFO survey, su	mmer - fall 1996

Figure 28: Percent maturity of female dogfish captured at the
three study locations in Atlantic Canada during the
summer - fall 1996

Figure 29: Number of progeny (NP) in relation to the total maternal
weight (TMW)(kg) for fecundity of the female dogfish
captured in the Minas Basin, during the N246 DFO
survey, and during the N249 DFO survey, summer - fall
1996

Figure 30:	Number of progeny(NP) in relation to the total maternal
	weight (TMW)(kg) for fecundity of female dogfish captured
	at the three study locations in Atlantic Canada during the
	summer - fall 1996

Figure 31:	Tag return locations for dogfish marked in the Minas Ba	isin
	during the summer of 1996	122

#### List of Appendices

Page	9
opendix 1: Raw data for female spiny dogfish captured in the	
Minas Basin from July 10 - October 10, 1996. TL	
= Total length; FL = Fork length	

ppendix 2: Raw data for male spiny dogfish captured in the	
Minas Basin from July 10 - October 10, 1996. TL	
= Total length; FL = Fork length	39

Appendix 3: Raw data for female spiny dogfish captured during the N246 Scotian Shelf DFO survey of July 1996. TL = Total length; FL = Fork length. . . . . 140

Appendix 5: Raw data for female spiny dogfish captured during	
the N249 Gulf of St. Lawrence DFO survey of	
September 1996. TL = Total length;	
FL = Fork length	49

Page
------

xv

Appendix 0. I daw data ior male spiriy dognari captured during
--

length	152
September 1996. TL = Total length; FL = Fork	
the N249 Gulf of St. Lawrence DFO survey of	

#### Acknowledgments

I would like to thank Ian Paterson and John MacMillan for their muscle power and practical ideas for fishing on a limited budget. My appreciation goes to Mark Johnston for illustrating the reproductive stages included in my thesis. Also to Bernie and boys of Delhaven, NS for allowing me onboard their fishing vessel and passing along some of their fishing knowledge. Thank you to DFO personnel, Tom Hurlbut and Jeff McRuer, for permitting me to participate on the annual groundfish surveys. My sanity would also like to thank: the gang on fifth floor in biology for conversations at the pool table, Wanda and Jeanette for listening, Donna and Ann for late night walks and coffee, and especially my family for listening to my biology babble in which they pretend to understand. And last but not the least, special thanks to Mike (Dinsey) Dadswell for his support and allowing me to work on a project that many people would avoid or consider a waste of time. Coffee time on fourth floor and Friday afternoon lunches at the Anvil will always be remembered.

#### Abstract

A total of 1684 dogfish were sampled to determine population characteristics from three regions in Atlantic Canada: the Minas Basin. the outer Bay of Fundy, and the Gulf of St. Lawrence. The posterior dorsal spine was removed from 211 male and 475 female dogfish for ageing. Ages were determined from the spines and annuli lost to wear were calculated using a spine dimension standardized equation. This study was the first attempt to define a complete reproductive stage assessment of a shark. A total of 475 females had the reproductive tracts removed and categorized as one of six stages. The Minas Basin sample exhibited the highest number of reproductively mature females (56.1%) and female dogfish captured in the outer Bay of Fundy were predominantly immature (97.6%). Maturity for 50% of female dogfish in Atlantic Canada was calculated at 17 years of age and a total length of 83.4 cm. Male dogfish age and total length were similar between the three regions but only 11 male dogfish were taken during the entire season in the Minas Basin (male:female ratio; 1:99). The Minas Basin female dogfish were significantly larger, older, and more mature than females in the other two sampling locations. Few comprehensive studies exist on spiny doafish in the Northwest Atlantic because it is has only recently been commercially exploited. If commercial exploitation of dogfish were to begin in Atlantic Canada it would be more economically and biologically successful as a bycatch fish with other fisheries.

#### Introduction

The spiny dogfish, Squalus acanthias Linnaeus, 1758, is a common shark of the North Atlantic and North Pacific Oceans. It occurs in the east and west coast of North America including the Bay of Fundy (Scott and Scott 1988). Comprehensive studies on its biology in the Bay of Fundy are non-existent, however, spiny dogfish is a slow growing, cartilaginous fish with a long lifespan. The Atlantic and Pacific populations are spatially separated and some of their characteristics have diverged enough to be considered separate species until recently (Jones and Geen 1976). Nammack et al. (1985) suggested that western Atlantic dogfish live approximately fifty years and reach average maximum lengths of 101 cm for females and 93 cm for males. Female dogfish of the North Pacific are known to live up to 95 years of age and reach lengths up to 124 cm (Ketchen 1975; Wood et al. 1979; Beamish and McFarlane 1985). These differences suggest Pacific dogfish may live longer and grow more slowly than dogfish of the western North Atlantic due to water temperature differences, but this possibility requires further study.

Squalus acanthias occurs in eastern Canadian inshore waters in the spring through fall period, appearing in the outer Bay of Fundy in May-June and departing in November-December (Scott and Scott 1988). Temperature may be the determining factor causing seasonal movements of dogfish. Studies indicate dogfish prefer offshore wintering grounds with temperatures between 6 °C to 11 °C (Jensen 1965).

Tagging studies by Templeman (1944; 1976; 1984), Holland (1957), and Jensen (1961; 1965) indicate dogfish school by size up to maturity and then by size and sex once they have matured. Templeman (1976) also recorded a transatlantic migration of dogfish from Newfoundland, Canada to Iceland.

Most studies have assumed spiny dogfish of the Northwest Atlantic are comprised of a single unit stock (Ford 1921; Templeman 1944; Templeman 1954; Shafer 1970; Nammack et al. 1985; Rago et al. 1994; Hurlbut et al. 1995). Preliminary work off the coast of North Carolina, USA indicates there were external differences between dogfish in the southern and northern range of the stock (Rulifson 1998). Annand and Beanlands (1986) completed protein electrophoresis on dogfish caught from the Gulf of Maine and the Scotian Shelf and found no difference between spiny dogfish from the two areas.

Male and female dogfish in the Atlantic reach sexual maturity at about 10 years of age and have different growth rates thereafter (Holden 1977). Males grow slower than females and are significantly smaller than females after maturity (Jensen 1965).

Spiny dogfish are ovoviviparous, meaning the eggs are hatched internally with no placental attachment to the mother (Holden and Meadows 1968; Ketchen 1972; Holden 1977; Jones and Geen 1977). Fetal development takes 18 - 22 months and from 1 - 25 pups are produced in each pregnancy. The sex ratio of pups is nearly 1:1 (Templeman 1944; Jones and Geen 1977a). There has been no actual data to confirm the season of the year in which mating occurs, but it has been suggested mating occurs between October and January, peaking some time in December (Ford 1921; Templeman 1944; Jensen 1965; Ketchen 1972). It is known mating occurs when water temperatures are low. Further investigation is required to determine the mating season of the spiny dogfish in the Northwest Atlantic.

Males are capable of mating each year and females every second year. Ovulation occurs just after mating and eggs average 4 cm in diameter. Fertilized eggs are enclosed in a capsule or 'candle' for 4 - 6 months, after which the embryos are released into the uterine cavity. Fecundity of the spiny dogfish increases with the size of the mother and varies from 1 to 25 pups (Templeman 1944; Holden and Meadows 1968; Ketchen 1972).

Age determination of all elasmobranchs is a difficult process because they do not possess calcareous otoliths and their scales are too small to determine annual ring formation. Reading annuli externally from the second dorsal spine is the most preferred method for determining the age composition of spiny dogfish populations (Holden and Meadows 1962; Ketchen 1975; Jones and Geen 1977a; Chilton and Beamish 1982; Soldat 1982; McFarlane and Beamish 1987).

The spine originates from the vertebral column, passes through the dorsal muscles and skin anterior to each dorsal fin. The portion embedded within the muscle is a cartilage rod, which keeps the spine in place. The spine consists of an outer enamel layer, a pigment layer, three layers of dentine, and a central

pulp cavity (Slauson 1982). The annulus forms because the dentine layers do not grow at the same rate as the upward growth of the spine. When spine growth is reduced, pigments are concentrated and the enamel layer thickens, producing an annulus (McFarlane and Beamish 1986). McFarlane and Beamish (1986) have identified these bands to be annual with oxytetracycline (OTC) injections from 18 recaptured dogfish. Only 1 out of the 18 recaptures did not show valid annuli with the OTC injections. Counting annuli is still very subjective and there are inconsistencies in ages due to the long lifespan and lack of adequate hard body parts.

The spiny dogfish is an opportunistic feeder whose diet includes fish, crustaceans, molluscs, and coelenterates (Templeman 1944; Holden 1965; Jones and Geen 1977b; Bonham 1984; Bowman 1984; Annand 1985; Nammack et al. 1985). Studies have shown 60 - 70% of the diet of dogfish over 60 cm in length were teleost fish. Their feeding habits reflect the abundance trends of certain fish species rather than preference (Bonham 1984). Consumption estimates and annual food intake of spiny dogfish by Bowman et al. (1984) suggest that dogfish predation may be increasing the mortality of commercially valuable species.

A large number of dogfish enter the inner Bay of Fundy during the summer fall (Dadswell et al. 1984a). These fish are relatively easy to capture and are considered a nuisance or undesirable species by fishermen and the public in Atlantic Canada (Templeman 1944; Salsbury 1986; Hurlbut et al. 1995). A preliminary study off Newfoundland in 1978 determined dogfish could not be commercially exploited in Canadian waters (Mercer et. al. 1979). In Great Britain and New Zealand, dogfish are a part of the local fish and chip market (Templeman 1944; Aasen 1964; Holden 1968; Salsbury 1986). Since 1990, the states of North Carolina, Virginia, Maryland, and Massachusetts, USA have developed a dogfish fishery, exporting mainly to the European markets. In January 1998, The National Marine Fisheries Service considered spiny dogfish over-harvested in the Northeast USA (NOAA, NMFS 1998).

Commercial fishermen in the USA are concerned that dogfish could become over-harvested before adequate biological data can be collected on this species (Jeff Gearhart pers. comm.). Northwest Atlantic spiny dogfish landings have increased from 519 metric tonnes (mt) in 1963 to 22 572 mt in 1993, with catches mostly from the United States and Canada. In the US, recorded landings have increased from under 500 mt in 1989 to over 20 000 mt annually since 1993 (Rago et al. 1994).

An experimental dogfish fishery was conducted in the fall of 1985 in the Scotia-Fundy region by the Department of Fisheries and Oceans. The experiment was designed to determine if small draggers could land economically feasible catches of quality dogfish. The groundfish bycatches exceeded the dogfish catches three to one and the idea to begin a directed dogfish fishery with drag gear in Atlantic Canada was ended (Salsbury 1986).

Comprehensive dogfish sampling in the inner Bay of Fundy and Minas Basin

has never before been attempted. The project will be of use to the Department of Fisheries and Oceans because it will provide information from a stationary sampling location on a non-commercial fish with no general annual assessment in Atlantic Canada.

The population characteristics of spiny dogfish, including: weight-length relationship, growth, reproductive state, percent maturity, and fecundity from morphological measurements and migration from tagging were studied. Data collected on groundfish surveys with the Department of Fisheries and Oceans in the southern Gulf of St. Lawrence N249 survey, the outer Bay of Fundy and its approaches N246 survey were compared to the Minas Basin samples. The final goal was to describe the basic population characteristics for this species and to determine if there were differences among dogfish in Atlantic Canada.

#### **Materials and Methods**

#### Location

The Minas Basin is located in the inner region of the Bay of Fundy along the province of Nova Scotia, Canada (Figure 1). The Bay of Fundy is known for the highest semi-diurnal tides in the world measured at maximum heights of 17.2 m off Burncoat Head of the Minas Basin (Larsen and Topinka 1985). The high tides are a result of the resonance between the oceanic tidal period and the natural period of the Bay of Fundy and the Gulf of Maine and the funnel effect created from the shape of the bay (Larsen and Topinka 1985).

All samples were taken and tags fixed between July 7, 1996 and October 10, 1996 using a 6.5 m research vessel. Dogfish were captured with either 22 mm longline hooks or gill nets of varying mesh sizes in the southern Bight of the Minas Basin between Blomidom Point and Boot Island (45.18N 65.42W) (Figure 1). One day of otter trawling with local fishermen was completed in Baxter's Harbour, NS (45.19N 65.30W) and shoreset gillnets were placed along Evangeline Beach (45.08N 65.41W) and Avonport Beach (45.06N 65.58W) to compare to other capture methods (Figure 1). Drift gillnets of varying mesh sizes were set off the research vessel throughout the season mainly for bait capture.

For comparison to the Minas Basin samples, special samples of dogfish were caught aboard the N246 and N249 groundfish surveys from July 4 - 16, 1996 and September 3 - 27, 1996. These surveys are conducted annually by the

Department of Fisheries and Oceans (DFO). Both surveys were aboard the 51m, RV Alfred Needler. Random set locations were selected prior to departure for both cruises according to DFO sampling protocols. The N246 survey was within the 4W and 4X NAFO divisions (Figure 2), along the eastern coast of Nova Scotia, approaches to the Bay of Fundy, and the Gulf of Maine. The N249 survey was conducted by DFO in the southern Gulf of St. Lawrence region of the 4T division (Figure 3).

#### **Capture Methods and Measurements**

All dogfish were measured for total length (TL) and fork length (FL) to the nearest 0.5 cm. Total length was determined as the length of the fish from the end of the snout to the tip of the upper lobe of the caudal fin when depressed to align with the longitudinal axis. Fork length was determined as the distance from the end of the snout to the central curve of the caudal fin. Weight was measured to the nearest 0.1 kg while at sea using a spring fish scale. The scale was calibrated using standardized one pound weights from laboratory balances. All posterior dorsal spines were removed from the sacrificed dogfish for ageing and if female, their reproductive tracts were removed and frozen for later analysis of the reproductive state.

A longline system was developed for sampling between Evangeline Beach and Blomidom Point, Minas Basin. The longline contained approximately 60, 25 mm hooks and soak time was 45 - 90 minutes. American shad (*Alosa*  sapidissima (Wilson, 1811)) caught in gillnets in the Minas Basin and in the Annapolis River, NS were cut up as bait for the hooks. Winter skate (*Raja ocellata* Mitchell, 1815) and dogfish were the only species caught using this method. Over 25% of the total number of marked dogfish were captured onboard a local flounder fishing vessel owned and operated by a local commercial fisherman from Delhaven, NS on August 28, 1996. Three separate tows were made using a 125 mm diamond otter trawl mesh, 24 m across within Baxter's Harbour, NS (Figure 1). The tows lasted 15 minutes, 30 minutes, and 10 minutes at 42 m depth. One Atlantic cod (*Gadus morhua* Linnaeus, 1758) was caught and released, and 12 winter flounder (*Pseudopleuronectes americanus* (Walbaum, 1792)) were kept as part of the fishermen's fishing quota.

Drift monofilament gillnets with stretch mesh sizes, 100 mm, 115 mm, 115 mm, and 125 mm were set off the research vessel in the Minas Basin for bait capture and as another method for capturing dogfish. Only American shad and 37 striped bass (*Morone saxatilis* (Walbaum, 1792)) were captured with this method. The shad were used as bait for the longline and the striped bass were returned to the water. Shoreset gillnets, the other method used to compare to the longline method, were not effective in capturing dogfish. The shoresets comprised of three monofilament gillnets with 115 mm, 125 mm, and 138 mm stretch mesh and were anchored during low tide off Evangeline Beach for one week and Avonport Beach, NS for two days. The nets were checked at each low tide for fish. The Evangeline shoresets did not capture any fish. Three dogfish

were captured during the two day set along Avonport Beach. This method was not effective enough in capturing dogfish to continue its use.

For the N246 and N249 DFO surveys a 20m otter trawl using 127 mm diamond mesh was towed behind the vessel for 30 minutes at each predesignated location. If the location was not adequate due to inappropriate physical characteristics (i.e.: rocky bottom) then a predesignated alternate station was used for the tow. For each tow 10 male and 10 female dogfish were set aside for sampling. All dogfish were measured for total length and fork length to the nearest centimetre, weight in kilograms to the nearest 0.01 kg using an electronic balance, the posterior dorsal spine was removed and the reproductive tract was removed from the females for later lab analysis.

#### Mark and Recapture Methods

All dogfish, in good condition, that were captured in the Minas Basin from July 7, 1996 to August 28, 1996 were measured for FL and TL to the nearest 0.5 cm, sex identified, marked with a tag, and released back into the water. A FT-1 Floy Dart tag was placed anterior to the first dorsal spine. Dogfish not in adequate condition to be returned to the water were sacrificed for further measurements, and reproductive tract and posterior dorsal spine removal.

Recapture location, date of recapture, and any other information given from the returned tags were recorded. The distance travelled from the Minas Basin tagging location to the recapture location was measured using a chartometer by the Kueffel and Esser Company. The chartometer is an instrument which measures the distance of a line using a tracing wheel calibrated to a graduated dial (Welch 1948). The wheel can be traced over a map to estimate the distance travelled along a pathway. The chartometer was calibrated using a 12 inch ruler. All measurements were taken in inches and converted to metric because the map was scaled in inches. Distances were traced along the coastline instead of a straight path since it was more likely the fish were travelling along the coastline than directly to the recapture location. These were only estimates of the actual distances travelled since it was not certain where the fish went between tagging and recapture.

#### Spine Removals and Ageing Techniques

The second dorsal spine was removed by placing the knife posterior to the spine. A cut was made parallel to the base of the spine down into the muscle tissue about 5 cm deep (Figure 4(a)). A second cut was anterior to the spine at a 45° angle was made into the tissue until reaching the first cut (Figure 4(b)). The spine was removed, placed into a labelled envelope, and frozen until later lab analysis. If the second dorsal spine was not adequate due to wear or damage, then the first dorsal spine was removed.

In the lab, the excess tissue around the base was excised and the spines were left to air dry for at least 24 hours. Amoural<sup>™</sup>, a vinyl cleaner normally used for car interiors and tires, was sprayed on the spines for further cleaning and to produce a shine which facilitated easier visualisation of annual ridges under the enamel layer. The spine grows in a series of dentine cones. Each cone represents one year with annual pigment bands forming between the enamel and dentine layers at the base of each cone (Slauson 1982; Figure 5).

Using Vernier calipers, all spines were measured for their spine length (SL), wear length (WL), spine base diameter (SBD), and wear point diameter (WPD) to the nearest 0.1mm (Figure 6). Under a light dissecting microscope, at 6.4 power and dark background, the annuli for each spine were counted from the spine base diameter to the wear point diameter (Ketchen 1975). There were problems counting the annuli on worn or damaged spines, as was the case in many other studies (Holden and Meadows 1962; Ketchen 1975; Chilton and Beamish 1982; Beamish and McFarlane 1985).

#### Calculations of Worn Spine Age and Growth

Annuli lost to wear were calculated by the method developed by Ketchen (1975). All spines with a wear diameter of 0 to 1.5 mm had their spine base diameter (mm) plotted against age (years). The diameter of 1.5 mm was determined as the spine base diameter at birth. Spines with wear points less than 1.5 mm were considered have no lost annuli. The curve of best fit was estimated from the equation:

Age =  $b(Spine base diameter)^{m}$  (Ketchen 1975)

where,

m = slope of the curve

b = y-intercept of the curve.

This equation was used to determine the number of years lost to wear from spine base diameters of worn spines. Two years were subtracted from this calculation before adding the number of lost annuli to the years counted from the spine base diameter to the wear point diameter because of the two year gestation period in which the spine would grow to about 1.5 mm in diameter. Ketchen's method was preferred for more accurate ages of dogfish because Nammack et al. 1985 found it better to compensate for lost annuli than rejecting worn spines.

The ages determined from Ketchen's method were used to obtain the von Bertalanffy growth relationship. The equation for the von Bertalanffy growth relationship is:

 $L_t = L_{\infty}(1 - e^{-K(t - t_0)})$ (Ricker 1975)

where,

 $L_t$  = total length (cm) at time t

 $L_{\infty}$  = theoretical maximum length in the population

t = age of the fish

 $t_0$  = a parameter for time when the total length equals zero

K = Brody growth coefficient

A computer generated curve fit of this equation was calculated from the observed data. The von Bertalanffy growth equation was fitted for dogfish

captured on the N246 survey, the N249 survey, and in the Minas Basin, for both males and females. The von Bertalanffy growth relationship was also determined for all males and all females captured during the summer-fail of 1996.

The weight-length relationship for males and females was obtained by plotting the weight (kg) against the total length (cm) of the fish. The equation for the weight-length relationship is:

log(weight) = a + b\*log(total length)(Ricker 1975)

where,

a = y-intercept of the regression line

b = regression coefficient.

The weight - total length relationship was plotted for the N246 survey, the N249 survey, and the Minas Basin for males and females. Overall, weight-length relationships for male and female dogfish was determined for the Atlantic Canada. The slope and standard error of each weight - total length relationship was compared between study locations to determine significant differences in growth. The equation to determine growth differences was:

 $t_{calculated} = \beta_1 - \beta_2$  (Steel and Torrie 1960)

 $S_1 - S_2$ 

where,

 $\beta_1$  = regression coefficient for a study location

 $\beta_2$  = regression coefficient for other study location

 $S_1$  = standard error for  $\beta_1$ 

 $S_2$  = standard error for  $\beta_2$ 

 $\sqrt{1}$  = square root of the entire denominator

The  $t_{calculated}$  value was compared to the  $t_{critical}$  value at alpha = 0.05 to determine if there were significant differences between the growth of male and female dogfish captured in the three study locations in Atlantic Canada (Steel and Torrie 1960).

#### **Reproductive Tract Removal and Measurements**

In all sacrificed females, whether immature or mature, the reproductive tracts were removed and placed in labelled bags for later analysis. The reproductive tracts consisted of the ovary, oviduct, and uterus. Using a knife or scissors, the ventral surface of the female was opened starting from the urinary pore and ending at the base of the pectoral fin (Figure 7).

Six stages of reproductive maturity and state in females were identified. The six stages are:

- Stage I Immature: no developing eggs, oviduct very thin and straight within the wall of the body cavity (Figure 8).
- Stage II Immature, with developing eggs: eggs having diameters > 20 mm, oviduct thin and straight within the wall of the body cavity (Figure 9).
- Stage III: Candle Stage: membrane surrounding embryos and their yolk sacs, very fragile, breaks open easily. Looks like a "candle". According to

the literature this stage represents the first 4 - 6 months after fertilization (Figure 10) (Templeman 1944; Holden and Meadows 1968; Ketchen 1972; Jones and Geen 1977).

- Stage IV: Post Candle Stage: Candle membrane no longer evident, embryos and attached yolks free-floating within the uterine cavity. Embryos measuring +16 mm in length, sex determination of embryos not possible, and developing eggs in the ovaries <20 mm in diameter (Figure 11).
- Stage V: Pup Stage: Can determine the sex of the individual pups. Pups look like small adults, and reach maximum total lengths up to 39.8 mm. Yolk sacs attached to the pups become smaller as they reach the end of the gestation period. Developing eggs in the ovaries >20 mm in diameter (Figure 12).
- Stage VI: Spent Stage: Uterine cavity flaccid. Eggs in the ovaries developed to 40 mm in diameter (Figure 13).

In the lab, the reproductive tracts were thawed and assigned to one of the six stages. The diameter of the eggs in the ovary were measured using Vernier callipers to the nearest 0.1 mm for reproductive tracts of Stages II to VI. The total length of embryos of Stage III and Stage IV were measured to the nearest 0.1 mm. The sex, total length (cm) and fork length (cm) to the nearest 0.5 cm, and weight (g) to the nearest 0.1g were determined for the pups in Stage V of reproductive development.

#### Fecundity, Maturity, and Embryo Development

Fecundity is defined as the number of progeny in the female before the next mating period (Caillet et. al. 1986). An estimation of fecundity was determined by plotting the number of progeny against the total weight (kg) of the mother to determine the curve of the form:

 $F = ax^{b}$  (Ricker 1975)

where,

F = fecundity

a = a constant

x = total maternal weight (kg)

b = an exponent, which is close to 1 when related to weight.

The curve was transformed into a straight line by logarithmic transformation:

 $\log(F) = (b)\log(x) + \log(a).$ 

Calculations of this line were developed for females captured from the N246 survey, the N249 survey, the Minas Basin, and for all three regions combined.

Mature female dogfish were considered fish with reproductive tracts described either as Stage III, Stage IV, Stage V, or Stage VI. Total maternal weight (kg) was plotted against the percent maturity of females within the weight parameter. A logistic curve fit was calculated to the data to determine the total number of the females at 50% maturity for the N246 region, the N249 region, the Minas Basin region, and all three areas combined.

The mean total length with the standard error for all embryos and pups within

Stage III, Stage IV, and Stage V was determined for each region and for all three areas combined. The mean egg diameter with their standard error for ovarian eggs in either Stage II, Stage III, Stage IV, Stage V, or Stage VI were determined for all separate study locations and the three locations combined.

# Analysis of Variance of Total Length, Age, and Maturity of Dogfish in the Three Study Locations

The median total length and median age of male and female dogfish were analysed for significant differences between the three study locations using the Kruskal-Wallis test by SigmaStat©(Tuerke et. al. 1993). The Kruskal-Wallis test is a non-parametric one way analysis test of variance on ranks. This nonparametric test was chosen over other tests because normality failed in the observed data and the population distributions were somewhat similar for all three locations. For this test the only assumption were the population distributions were continuous and the same shape (Farrell 1997; Netter and Wasserman 1974). The total length and age distributions for both male and female dogfish were mostly skewed left.

Three categories determined the percent maturity of female dogfish dependent on the maternal weight. The ranges for the maternal weight were: 0.5 - 1.49 kg, 1.5 - 2.49 kg, and  $\geq$  2.5 kg. These values were then analysed using the Kruskal-Wallis rank test to determine significant differences among the three maternal weight ranges of the three separate study locations.

#### **Estimation of Growth Parameters**

All computer generated curve fits were calculated using Sigma Plot© by Jandel Scientific (Tuerke et al. 1993). Jandel Scientific uses the Marquardt-Levenberg algorithm to find the coefficients for the best fit of the equation to the actual data. The algorithm verifies the parameters to minimize the sum of the square differences between the observed and predicted values of the data.

All data for males and females were separated because of the difference in size after maturity. Calculations were also determined separately for dogfish capture on the N246 survey, the N249 survey, and in the Minas Basin.

#### Results

#### **General Population Characteristics**

A total of 1115 spiny dogfish were captured off the research vessel in the Minas Basin, from July 7, 1996 to October 10, 1996 (Table 1). Of these captured dogfish, 751 were marked and released, and 364 were sacrificed for reproductive tract and spine removal. An additional 244 dogfish were captured aboard a local flounder fishing vessel in Baxter's Harbour, NS on August 28, 1996, marked and released (Figure 1). A total of 216 spiny dogfish were sampled during the N246 DFO survey and 114 during the N249 DFO survey at various set locations (Figure 14; Figure 15).

The ratio of males to females captured at each of the three regions differed (Figure 16; Figure 17). On the N246 DFO survey, 1:0.7 males to females were captured. On the N249 DFO survey, 1:1.5 males to females were captured. The Minas Basin dogfish collections were predominantly female. A total of 1.0:99.2 males to females were captured at this location.

Female dogfish ranged in total lengths from 51 - 87 cm on the N246 survey, from 64 - 101 cm on the N249 survey, and from 64 - 113 cm in the Minas Basin. Male dogfish ranged in total lengths from 53 - 86 cm on the N246 survey, from 64 - 87 cm on the N249 survey, and from 74 - 86 cm in the Minas Basin (Figure 16; Figure 17; Table 2). The age range for female dogfish captured on the N246 survey was 3 - 36 years, on the N249 survey from 9 - 27 years, and in the Minas
Basin from 10 - 34 years. Male dogfish ranged from ages 2 - 29 years on the N246 survey, from 9 - 24 years on the N249 survey, and from 10 - 21 years in the Minas Basin (Figure 18; Figure 19; Table 2). The age range was much wider for both male and female dogfish captured on the N246 survey compared to the other two locations. Younger dogfish were captured on the N246 survey compared to the normared to dogfish captured in the Minas Basin and on the N249 survey.

Normality failed in all comparisons (Female TL: p < 0.0001; Male TL: p < 0.0001; Female Age: p < 0.0002; Male Age: p = 0.0036) (Table 3). Total lengths and ages of females varied significantly among the three locations (Female TL: p < 0.0001; Female Age: p < 0.0001; Male Age: p = 0.0217). The total lengths of males were significantly different (p < 0.0001) between the three regions but the ages were not (p = 0.5038).

## Weight-Length Relationship and Growth Comparisons

The weight-length relationships of female spiny dogfish for the three study locations were similar (Figure 20) as they were for male spiny dogfish (Figure 21). The combined weight-length relationship for female spiny dogfish from Atlantic Canada was:

 $log(W) = 3.5log(TL) - 6.1; r^2 = 0.94$ 

and for male dogfish was:

 $log(W) = 2.8log(TL) - 5.1; r^2 = 0.93$ 

where, weight was measured in kilograms and total length in centimetres (Figure

Female dogfish varied more in weight than males at similar lengths. The Minas Basin female dogfish varied more in their weight-length relationship than the N249 or N246 female dogfish. The b-values ranged from 3.2 - 3.4 meaning they exhibited allometric growth, therefore they were growing slightly more in weight than they were in length (Ricker 1975). Male spiny dogfish did not have as high b-values (between 1.6 - 2.9) therefore weight was not increasing as much as length.

Growth, using the weight-length relationship, was equal for male and female dogfish among the three regions, except the between male dogfish from the N246 DFO survey and the Minas Basin ( $t_{calculated} = 2.2580$ ; Table 4). This comparison may not be valid due to the small sample size of male dogfish captured in the Minas Basin and not directly related to differences in growth between males captured in the two regions.

### Age and Growth

A total of 644 spines were analyzed during this study: 344 from the Minas Basin, 200 from the N246 DFO survey, and 100 from the N249 DFO survey. The spine base diameters (SBD) of spines  $\leq$  1.5 mm in diameter were standardized for age (years) to determine the equation of best fit for annuli loss (Figure 23). This equation was used to determine the number of annuli lost to wear in older spines (Ketchen 1975). Male dogfish have an equation for annuli loss of:

22).

Age =  $1.4(SBD)^{1.5}$ ;  $r^2 = 0.49$ .

The female dogfish equation for annuli loss was determined as:

Age =  $1.8(SBD)^{1.3}$ ;  $r^2 = 0.52$ .

The spine base diameter for all wear points greater than 1.5 mm were standardized into either of these equations depending on the sex of the individual to determine the number of lost annuli. The calculated number of lost annuli were added to the ages determined on the spines' external readings from the SBD to the point of wear. The new ages were used to determine the von Bertalanffy growth equation for male and female dogfish at all three study locations and the study locations combined (Figure 24; Figure 25; Figure 26; Table 5).

The asymptotic length,  $L_{\infty}$ , was smallest for both male ( $L_{\infty} = 77$ ) and female ( $L_{\infty} = 80$ ) dogfish captured during the N246 DFO survey and greatest in the Minas Basin (male  $L_{\infty} = 81$ ; female  $L_{\infty} = 97$ ). The birth size,  $t_0$ , was close to 2 for most von Bertalanffy growth equations, ranging from 1.9 - 2.8 for female dogfish and from 1.7 - 2.13 for male dogfish in the three regions. The Brody growth coefficient, K, was between 0.14 and 0.20 for female dogfish in the three regions and between 0.22 and 0.27 for males captured in the three regions. A larger K for males indicates faster growth for male dogfish than for female dogfish.

For all three regions combined, the von Bertalanffy growth equation for female dogfish was:

 $L_t = 91(1 - e^{-0.17(t-2.5)})$ 

and for male dogfish captured in all three study locations:

 $L_t = 78(1 - e^{-0.25(t-1.8)})$ (Figure 26).

#### Maturity

Female dogfish captured in the Minas Basin were found to be in all stages of development with 56% of the total sample reproductively mature (Table 6). Female dogfish captured during the N246 DFO survey were almost completely immature (97.6%). Female dogfish from the N249 survey showed all but Stage II of reproductive development and 72% were immature.

In the Minas Basin, most of the mature females were captured in the early part of the run from July to late August. The majority of the immature females in the Minas Basin were captured from September to October. Most of the males captured in the Minas Basin were not taken until October.

A frequency distribution of maternal fish weight (kg) to percent maturity determined the weight of female dogfish at 50% maturity for the three study regions and the regions combined (Figure 27; Figure 28). Fifty percent maturity of the sampled female dogfish on the N246 and N249 DFO surveys were both 2.9 kg weight or 90.6 cm and 85.1 cm total length. Age at 50% maturity for female dogfish on the N246 survey was 26 years and for female dogfish on the N249 survey 17 years. Female dogfish from the N246 DFO survey were mostly immature (97.6%) (Table 6), therefore age at 50% maturity was overestimated probably from lack of mature females in the sample. In the Minas Basin 50%

maturity was calculated to be at a weight of 2.7 kg, total length of 82.4 cm, and age of 16 years. For all three study locations combined 50% maturity was reached at a weight of 2.7 kg or total length of 83.4 cm. The mean age of female dogfish at 50% maturity in Atlantic Canada was about 17 years.

The percent maturity in relation to maternal total weight between the N246 DFO survey, the N249 DFO survey, and the Minas Basin collections were categorized into three maternal total weight categories: from 0 - 1.49 kg, 1.5 - 2.49 kg, and 2.5 - all above weights. There were no differences between the 0 - 1.49 kg female weight category for the three study locations because samples were all immature in this weight range. There were no statistically significant differences between the maturity of females within the 1.5 - 2.49 kg weight range in all three study locations (Kruskal-Wallis; p = 0.2124). Female dogfish categorized  $\geq$  2.5 kg in total weight, showed a statistically significant difference between the median % maturity values for the three study locations (Kruskal-Wallis; p = 0.0338). This indicated that maturity was different among the regions in Atlantic Canada. Female dogfish in the Minas Basin were more mature than female dogfish captured in the outer Bay of Fundy and Gulf of St. Lawrence.

## Fecundity and Reproductive Development

Litter sizes ranged from 1 - 11 embryos, with an average of 5.23 embryos per mature female (Figure 30). There was variability in the number of embryos per female of any given weight at all study locations. Minas Basin females had the most variable litters (Figure 29). Fecundity was represented for the N246 DFO survey, the N249 DFO survey, and the Minas Basin combined as:

log(Number of progeny) = 0.8log(Total maternal weight) + 0.3

Embryonic lengths measured from 16.1 - 398.0 mm in total length, with a mean total length of 39.8 mm for Stage III embryos, 72.3 mm for Stage IV embryos, and 224.8 mm for Stage V pups (Table 7). A total of 232 pups were male and 235 pups were female. There was no significant difference between size of male or female pups in Stage V development (Mann-Whitney; p = 0.3137).

#### Migration

A total of 17 recaptures were obtained from 995 spiny dogfish tagged during 1996 in the Minas Basin with an average of 276 days at large and an average distance traveled of 569 km (Figure 31; Table 8). Recaptures were 0.02% of the total number of spiny dogfish marked and released in the Minas Basin. Most recaptures were returned just after the opening of a local fishery.

Four recaptures were returned off Yarmouth, NS at the beginning of the winter lobster fishery started in December (#1694, #1190, #1646) and after the beginning of the April lobster season opening (#1580 and #1294). The Sandy Cove, NS tag (#1640) was captured in a herring weir and one off Port Maitland, NS (#1274) was captured by handline.

The Rhode Island and Cape Cod tag returns (#1264, #1053, and #1702)

were captured just after the local dogfish fishery opened and another tag (# 212) from the same area was captured by rod and reel. Tag #212 traveled the longest distance, 1056 km, in only 144 days. This dogfish traveled an average of 7.3 km per day. The Gloucester, Massachusetts tag returns (#1226, #1904, and #1901) were captured just after the opening of their local dogfish fishery. Only two tags were returned from the Minas Basin close to the tagging location (#1425) on May 26, 1997 and the other in the Avon River on August 6, 1997 (#1480). The tag at liberty for the longest time (#1159) was captured off Grand Manan, NB, 630 days after release. Tag #1264 was captured on the same day twice. First by a NOAA groundfish survey vessel and later by a local commercial fisherman off Cape Cod.

#### Discussion

#### **General Population Characteristics**

The Bay of Fundy is a unique ecological system and provides an important link in the life history processes of many fish species along the eastern North American coastline (Dadswell et al. 1984a). The inner reaches of the Bay of Fundy are warmer in the summer than other localities on the Canadian east coast and may be uilized by fish with specific life history needs (Bousefield and Liem 1958). Spiny dogfish have never before been studied in the Bay of Fundy, or compared to other regions in Atlantic Canada. This study has indicated variations exist among spiny dogfish aggregations at different geographical regions in Atlantic Canada and these differences may be due to life history strategies of the local dogfish stock.

Female dogfish captured in the Minas Basin were larger and more mature than females captured during the Scotian Shelf survey or during the Gulf of St. Lawrence survey. Spiny dogfish caught during the N246 DFO survey were taken mostly in the outer approaches of the Bay of Fundy, very few were present along the Nova Scotian shelf or the offshore region at that time of the year. Female spiny dogfish captured in the Gulf of St. Lawrence were somewhat similar in their size range to females in the Minas Basin, but were not as reproductively mature.

There was a 99:1 ratio of females to males in the Minas Basin. Mature

females were most predominant at the early and middle periods of the run in the Minas Basin and as the season progressed, immature females and males became more common. Mature females may have been entering the Minas Basin, seeking the warmer waters found there during the summer, to assist internal growth of their offspring (Bousefield and Liem 1958). Spiny dogfish have one of the longest gestation periods of any vertebrate species, from 22 - 25 months (Ford 1921; Templeman, 1944; Jensen 1965). Other studies have indicated that mature dogfish tend to be closer inshore, which is generally a warmer region of the ocean, than immature dogfish (Jensen 1965; Nammack et al. 1985; Hurlbut et al. 1995). The large number of female dogfish in the Minas Basin indicates that the Bay of Fundy may be a significant location in Atlantic Canada for the reproductive success of this species.

#### Growth

Compared to two other studies in the Northwest Atlantic (Slauson 1982; Nammack et al. 1985) the maximum asymptotic length ( $L_{\infty}$ ) for all three locations obtained during this study were lower and the growth coefficient (K) was usually higher. The highest  $L_{\infty}$  was 97 cm for female dogfish in the Minas Basin, whereas Slauson (1975) and Nammack et al. (1985) calculated an  $L_{\infty}$  greater than 100 cm for female dogfish off Massachusetts and Rhode Island. The K values were also lower for females in these other studies at 0.07 and 0.09, therefore growing slower, as compared to 0.14 and 0.20 for female dogfish captured in the Minas Basin and during the N246 DFO survey. From this information, it can be assumed dogfish in the northern range had a lower mean asymptotic total length but grew at a faster rate than the southern relatives.

Nammack et al. (1985) and Slauson (1982) both found negative  $t_0$  values for their von Bertalanffy equations of dogfish growth but in this study  $t_0$  was positive. Their studies did not account for the gestation period and  $t_0$  began at birth instead of at conception. It is more appropriate to consider growth beginning after conception in the uterus, and embryo lengths can be incorporated in to the von Bertalanfy growth equation.

Lower L<sub>∞</sub> than the observed lengths may indicate my sample size was not large enough at the higher end of the growth range. Also, L<sub>∞</sub> could have been acting as only the mean maximum size for the entire population. Larger individuals would be expected in an unexploited stock above this average size (Nammack et al. 1985). Males have a greater K value than females, therefore males reach their theoretical maximum length (L<sub>∞</sub>) sooner and grow faster than females. From my growth curve estimations the calculated time when fish length equals zero ( $t_0$ ) for males and females was close to 2 years which is equivalent to the time pups grow before birth within the mother.

The weight-length relationship of male and female dogfish are similar up to about 50 cm total length and then greater for females after this length (Nammack et al. 1985). Male dogfish tend to grow isometrically throughout their entire lifetime. Female dogfish grow isometrically up to maturity and show mostly allometric growth after maturity. Females tend to grow larger and vary more in size after maturity because of pregnancy. The number of pups per pregnancy increases with the age of the female dogfish (Templeman 1944; Jensen 1965; Holden 1977; Jone and Geen 1977a).

# Ageing

Accurate age determination of fishes is an important requirement in assessing and achieving proper management strategies for any commercially exploited species. It is helpful to know the size and age at which a species become sexually mature so fishing can be restricted and reproduction can occur before being exposed to harvesting pressure. All elasmobranchs are a difficult group for determining age, growth, and reproduction because they are elusive, have minimal commercial value, and possess few anatomical structures for accurate age verification.

Age determination using the second dorsal spine of the spiny dogfish has been utilized in other studies (Kaganovskaia 1933; Templeman 1944; Aasen 1964; Holden and Meadows 1962; Ketchen 1975; Soldat 1982). Spiny dogfish are one of the few elasmobranch species in which dorsal spine circuli have been directly validated as annual, but only for the Canadian west coast stock (Beamish and McFarlane 1985). Age validation has not been conclusively shown for spiny dogfish of the Northwest Atlantic.

This study found spiny dogfish captured in Atlantic Canada ranged in age

31

from 2 - 36 years of age. Maximum ages were younger than anticipated from the literature. Other studies in the Northwest Atlantic have suggested that spiny dogfish can live up to 50 years of age but have only demonstrated age counts up to 35 years (Slauson 1982; Nammack et al. 1985). Jones and Geen (1977a) noted Ketchen's technique (1975) of compensating for missing annuli of worn tips produces better results than rejecting worn spines (Nammack et al. 1985). Variations in length at age is attributed to the difficulty in interpreting rings along the spine and individual differences among the populations.

One method not examined during this study which is often used for mammals is measuring the weight of the eye lens. The eye lens is an ectodermal structure which grows throughout the lifetime of the organism and its mass should be proportionate to the length and age of the organism (Friend 1968). Due to the lens position in the body it does not undergo wear like many other structures used for age determination (Morris 1972). Siezen (1989) has determined a relationship between lens weight and age of spiny dogfish from measurements of lens weight, body length, and spine base diameter. This method has also proven effective in ageing smooth dogfish, *Mustelus canis* (Mitchill, 1815) (Zigman and Yulo 1979). Eye lens verification in dogfish could be used to increase the accuracy of the spine technique and create a standardized ageing technique for all elasmobranchs.

### **Reproduction and Maturity**

It is important to develop stages for the gestation period in order to determine the timing for conception, and for assessing more accurately, recruitment into the dogfish population. This is a first attempt to determine a general reproductive assessment in stages for a shark species. Since most female dogfish captured in the Minas Basin were in Stage III and Stage V of reproductive development this would indicate that mating occurred between November to March, following Jensen's (1965) observations. The mating season, if there is one, for spiny dogfish is still relatively uncertain. Male spiny dogfish were more prevalent in offshore areas so it would be reasonable to suggest mating occurs offshore (Ford 1921; Templeman 1944; Jensen 1965; Slauson 1982).

Embryonic growth and development in Atlantic Canada was similar to other studies in the Northwest Atlantic (Ford 1921; Templeman 1944; Slauson 1982). Pupping or the birth of the young may be seasonal. Female dogfish in Stage III and Stage V in Atlantic Canada during the summer, estimates conception occurred no later than February. Evidence of spent females later in the season (October) could indicate the Minas Basin as a pupping ground for this species. It can be suggested from this study the pupping season for females coming to Atlantic Canada occurs anywhere between October to February.

Dogfish from the Pacific have had an estimated 50% maturity at 29 - 31 years of age (Ketchen 1972; Jones and Geen 1977). Templeman (1944) suggested that 50% maturity was reached at 12 years of age for dogfish

33

captured off Newfoundland. Maturity for 50% of the female dogfish was estimated between 16 - 26 years of age by this study. Dogfish in the Northwest Atlantic appear to be maturing at a faster rate than the west coast stock. Variation in the age of 50% maturity is likely due to changes in environmental conditions (Hanchet 1988; Saunders and McFarlane 1993). It has also been suggested that most fish species do not mature at a certain size or age but along a certain range of ages and lengths, dependent upon the level of environmental stress encountered (Stearns and Crandall 1984). These observations may also be appropriate for spiny dogfish.

Spiny dogfish have been known to vary in maturity and fecundity within small geographical areas, like the west coast of Canada (Ketchen 1972), the Northeast Atlantic (Holden and Meadows 1968), and even in the southern hemisphere (Hanchet 1988). Environmental stress could be considered variation in food supply, constant fluctuations in water temperature, storm activity, or even fishing pressure on other species in which dogfish prey upon. Therefore, dogfish under environmental stress could have slower growth and reach maturity later in one part of the population or have faster growth and earlier maturity in another part of the population. The Northwest Atlantic is known to be one of the most environmentally fluctuating part of the world oceans (Hildebrand 1984), therefore spiny dogfish aggregations can vary within Atlantic Canada, as shown in this study.

## Management of Spiny Dogfish

Most of the literature suggests spiny dogfish comprise of one unit stock in the Northwest Atlantic (Ford 1921; Templeman 1944; Shafer 1970; Nammack et al. 1985; Rago et al. 1994; Hurlbut 1995), but no accurate genetic research has been completed to verify this assumption. Evidence indicates that dogfish in different regions have unique fecundity and growth rates, and migration depends on the maturity of the fish (Nammack et al. 1985; Anderson 1990). Spiny dogfish are known to have at least two separate stocks in waters off British Columbia in Georgia Strait and Hectate Strait (Ketchen 1975). Three separate stocks existed in British waters known as the Scottish-Norwegian Stock, the Channel Stock, and the Atlantic Stock (Holden 1965).

In my study there were two defined groups traveling from the Minas Basin. One group moved and overwintered in the area off the southeast coast of Nova Scotia and another continued south to Massachusetts in the Cape Cod region along the coastline. It can be suggested that there may be more than one unit stock in Atlantic Canada. Preliminary observations of dogfish captured off the coast of North Carolina have indicated some morphological differences to the northern relatives, and there may be northern and southern stocks (Rulifson 1998). Fisherman from North Carolina also believe there is an aggregation of spiny dogfish in the deeper coastal waters that remains year round and the group is a nuisance to fisherman after other commercial species in these waters (Roger Rulifson pers. comm.). Accepting there is more than one unit stock would change the approach to management of this species in a sustainable harvest level along the Atlantic coast.

Annual DFO survey results in Atlantic Canada have shown an increase in spiny dogfish abundance since 1987 (Hurlbut et al. 1995). Inshore fishermen regard spiny dogfish as a nuisance, that interferes with fishing operations more than any other species. Spiny dogfish abundance has been so great to temporarily stop fishing activities in areas of the Bay of Fundy (Bernard Millett pers. comm.). The ecological role of spiny dogfish on other species has not been assessed. It has been suggested that the increase in spiny dogfish abundance is the reason for the decline of many commercially valuable species in Atlantic Canada. A directed fishery for spiny dogfish to counteract this supposed decline of other fish species on a large enough scale in Atlantic Canada has been suggested but never implemented (Mercer 1979; Hurlbut et al. 1995).

A large scale directed dogfish fishery along the east coast of the United States for the past eight years has become a lucrative industry with most of the catch exported to Europe and Asia (NOAA, NMFS 1998). The markets demand large dogfish, and catches have predominantly consisted of pregnant females. Since the Minas Basin dogfish stock is comprised mainly of mature females it would be an ideal economical and efficient target location to begin a directed dogfish fishery for the current world market. It may not be economically feasible to create a large scale directed dogfish fishery in Atlantic Canada, but it could become a bycatch fishery. Dogfish captured during other directed commercial

36

fishery operations could be brought in to market as a bycatch incidence fish to reduce its abundance and nuisance to fishermen. This study will provide critical biological information for management of the spiny dogfish stock granted it becomes a commercially exploited species in Atlantic Canada.

Spiny dogfish abundance should not be attributed to the decline of other commercial species such as cod. Dogfish have been indicated in many papers as the cause for declining stocks of other species (Salsbury 1986; Rago et al. 1994; Hurlbut 1995). Dogfish are opportunistic and eat many different species without distinct preferences. A study just released from the Northeast Fisheries Science Center of Woodshole, Massachusetts, spring and autumn trawl surveys from 1993 - 1997, contains information about the stomach contents of spiny dogfish. Of 8400 stomachs sampled during the spring and autumn surveys, only 14% contained fish species (NOAA, NMFS 1998). Ctenophores and crustaceans represented the most frequent species identified in the stomachs of doafish. The surveys indicated a majority (40 - 46%) of the dogfish stomachs were empty probably due to regurgitation. Food contents in dogfish stomach were not scientifically identified in this study, but it was noted that many were full of crabs common to the Minas Basin. In areas where dogfish are abundant and regarded as a nuisance, they should not be reduced in biomass solely on the basis that they may predate on more commercially acceptable species. More accurate information on the feeding habits of spiny dogfish should be obtained before a fishery is established to determine the ecological role of spiny dogfish in food

37

web.

The abundance of spiny dogfish of the Northwest Atlantic stock has increased since the 1960s (Rago et al. 1994; Hurlbut 1995). In 1977, The Magnuson Act was established and landings decreased significantly due to the exclusion of foreign vessels in US waters. With the beginning of a dogfish fishery in Massachussets, Virginia, and North Carolina in 1990. landings have since increased by a factor of four. Also, the number of discards have increased and depending on the percentage of the discards which die, then the level of the current fishing mortality could be much higher than recorded. The total catch in 1993 was 36 000 mt but may have been as high as 50 000 mt (Rago et al. 1994). The spiny dogfish population of the Northwest Atlantic has probably declined in abundance due to increased exploitation (Hurlbut et al. 1994; Rago et al. 1994; NOAA, NMFS 1998). Length frequencies from both USA commercial landings and research vessel surveys indicate a decline in the average length of females (Rago et al. 1994). The median weight for females landed since the beginning of this fishery has dropped by 1.5 kg (NOAA, NMFS 1998).

Spiny dogfish are sensitive to overfishing due to their life history patterns. Stock and recruitment are directly related and litter sizes may increase to a point as stock density decreases (Holden and Meadows 1968). Uncertainty exists on the data and parameters used in assessing and managing the Northwest Atlantic spiny dogfish stock. More accurate information is needed on total catch estimates with both catch and discard data, and more sex and length characterization of the total catch (Rago et al. 1994).

This past winter spiny dogfish were considereded an overharvested species by the National Marine Fisheries Service in the United States with little information from the southern end of the range to justify this declaration (NOAA, NMFS 1998). This study will provide a basic overview of the life history of spiny dogfish in Atlantic Canada and provide information at the northern end of the range for the stock.

A management program with appropriate targets for stock biomass and fishing mortality should be established. The model developed by Wood et al. (1979) for the British Columbia stock was the only age structured model applied successfully to elasmobranchs. The model was initially used for marine mammals and due to similarities in life history patterns, it could be used for elasmobranchs. Wood et al. (1979) concluded from the model that mortality is a a density -dependent factor of the changes in the stock caused from exploitation. It has been shown in whale stocks that recruitment changes greatly and influences sustainable yields at different stock levels. Further analysis is necessary to determine if change in stock levels influence similar biological parameters in sharks, such as spiny dogfish, as they do in whale stocks (Anderson 1990).

# Bibliography

- Aasen, O. 1964. The exploitation of the spiny dogfish (Squalus acanthias L.) in European waters. Fiskeridir. Skr-fisk. 13: 5 16.
- Anderson, E. D. 1990. Fishery models as applied to Elasmobranch fisheries. NOAA Tech. Rep. NMFS 90: 473 - 484
- Annand, C. and D. Beanlands. 1986. A genetic stock study of dogfish in the Northwest Atlantic. NAFO SCR Doc. 86/102. N1229. 5 pp.
- Beamish, R. and G. A. McFarlane. 1985. Annulus development of the second dorsal spine of the spiny dogfish (*Squalus acanthias*) and its validity for age determination. Can. J. Fish. Aquat. Sci. 42: 1799 1805.
- Bonham, K. 1984. Food of the dogfish, *Squalus acanthias.* Wash. Dept. Fish. Wildl. Serv. Fish Res. Pap. 1: 25 36.
- Bousefield, E. L. and Leim, A. H. 1958. The fauna of the Minas Basin and the Minas Channel. National Museum of Canada Bulletin. 166: 1 30.

- Bowman, R., R. Eppi, and H. Grosslein. 1984. Diet and consumption of spiny dogfish in the Northwest Atlantic. Int. Counc. Explor. Sea. Woodshole, Massachusetts.
- Caillet, G. M., M. S. Love, A. W. Ebeling. 1986. Fishes. A Field Guide and
  Laboratory Manual on Their Structure, Identification, and Natural History.
  Wadsworth Publishing Company. Belmont, California. 194 pp.
- Chilton, D. E. and R. J. Beamish. 1982. Age determination methods forfishes studied by the groundfish program at the Pacific biological station.Can. Spec. Pub. Fish. Aqua. Sci. 60: 94 98.
- Dadswell, M.J., R. Bradford, A. H. Leim, D. J. Scarratt, G. D. Melvin, and R, G, Appy. 1984a. A review of research on fishes and fisheries in the Bay of Fundy between 1976 and 1983 with particular reference to its upper reaches. Can. Tech. Rep. Fish. Aqua. Sci. 1256: 163 - 294.

Farrell, P. 1997. Statistical Consulting Center. Acadia University. Wolfville, NS.

Ford, E. 1921. A contribution to our knowledge of the lif histories of the dogfishes landed at Plymouth. J. Mar. Biol. Ass. U.K. 12: 468 - 505.

- Friend, M. 1968. The lens technique. Trans. 33rd. N. Am. Wildl. Nat. Res. Conf. 33: 279 298.
- Hanchet, S. 1988. Reproductive biology of *Squalus acanthias* from the east coast, South Island, new Zealand. New Zeal. J. Mar. Fres. Res. 22: 537 549.
- Hildebrand, L. P. 1984. Oceanographic setting. In: R. C. H. Wilson and R. F.
  Addison (eds.). Department of the Environment/ Department of Fisheries and
  Oceans/ Department of Energy. Mines and Resources. Canada. 8 30.
- Holden, M. J. 1965. The stock of Spurdogs (*Squalus acanthias* L.) in British waters and their migrations. Fish. Invest. Ser. 2. 24: 1 -19.
- Holden, M. J. 1968. The rational exploitation of the Scottish-Norwegian stock of spurdogs (*Squalus acanthias* L.). Fish. Invest. Ser. 2. 25: 1 27.
- Holden, M. J. 1977. Elasmobranchs. In: J. A. Guillard (ed.). *Fish Population Dynamics*. J. Wiley and Sons Ltd., London. 187 214.

- Holden, M. J. and P. S. Meadows. 1962. The structure of the spine of the spur
  dogfish (*Squalus acanthias* L.) and its use for age determination. J. Mar. Biol.
  Assoc. 42: 179 197.
- Holden, M. J. and P. S. Meadows. 1968. The fecundity of the spurdog (Squalus acanthias L.). J. Cons. Perm. Int. Explor. Mer. 28: 418 424.
- Holland, G. A. 1957. Migration and growth of the dogfish shark, *Squalus acanthias* (Linnaeus), of the Eastern North Pacific. Wash. Dept. Fish. 2: 43 59.
- Hurlbut, T., G. Neilsen, R. Hebert, and D. Gillis. 1995. The status of spiny dogfish (*Squalus acanthias*, Linnaeus) in the southern Gulf of St. Lawrence (NAFO Division 4T). DFO Atl. Fish.Res. Doc. 95/42. 38 pp.
- Jensen, A. C. 1961. Recaptures of tagged spiny dogfish, *Squalus acanthias*. Copeia. 228 - 229.

Jensen, A. C. 1965. Life history of the spiny dogfish. U. S. Fish. Wild. Ser. Fish. Bull. 65: 527 - 543.

- Jones, B. C. and G. H. Geen. 1976. Taxonomic reevaluation of the spiny dogfish (*Squalus acanthias* L.) in the northeastern Pacific ocean. J. Fish. Res. Bd. Can. 33: 2500 2506.
- Jones, B. C. and G. H. Geen. 1977. Reproduction and embryonic development of spiny dogfish (*Squalus acanthias*) in the Strait of Georgia, British Columbia. J. Fish.Res. Bd. Can. 34: 1286 1292.
- Jones, B. C. and G. H. Geen. 1977a. Age determination of an Elasmobranch (*Squalus acanthias* L.) by x-ray spectrometry. J. Mar. Biol. Assoc. 42: 179 - 197.
- Jones, B. C. and G. H. Geen. 1977b. Food and feeding of spiny dogfish (*Squalus acanthias*) in British Columbia waters. J. Fish. Res. Bd. Can. 34: 2067 -2078.
- Kaganovskaia, S. 1933. A method of determining the age and the compsition of the catches of the spiny dogfish (*Squalus acanthias* L.). J. Fish. Res. Bd. Can Trans. Ser. 281: 139 141.

- Ketchen, K. S. 1972. Size at maturity, fecundity, and embryonic growth of the spiny dogfish (*Squalus acanthias*) in British Columbia waters. J. Fish. Res.
  Bd. Can. 29: 1717 1723.
- Ketchen, K. S. 1975. Age and growth of dogfish *Squalus acanthias* in British Columbia waters. J. Fish. Res. Bd. Can. 32: 43 - 59.
- Larson, P. F. and J. A. Topinka. 1985. Fundy Tidal Power Development: Preliminary Evaluation of its Environmental Consequences to Maine. A Report to the Maine State Planning Office by Bigelow Laboratory for Ocean Sciences. A Division of the Northeastern Research Foundation, Inc. Tech Rep. 35.
- McFarlane, G. A. and R. J. Beamish. 1986. A tag suitable for assessing longterm movements of the spiny dogfish and preliminary results from use of this tag. N. Am. J. Fish. Manag. 6: 69 - 76.
- McFarlane, G. A. and R. J. Beamish. 1987. Validation of the dorsal spine method of age determination for spiny dogfish. In: Age and Growth of Fish. The Iowa State University Press, Iowa.

- Mercer, K. M., R. J. Sesk, and T. Fosnes. 1979. Development of the dogfish fishery in Newfoundland during 1978. Government of Newfoundland and Labrador. Dept. Fish. Dev. Bran. Rep. 4.
- Morris, P. 1972. A review for mammalian age determination methods. Mam. Rev. 2: 69 104.
- Nammack, M. F., J. A. Musick, and J. A. Colvocoresses. 1985. Life history of the spiny dogfish off the Northeastern United states. J. Am. Fish. Soc. 114: 367 - 376.
- Netter, J. and W. Wasserman. 1974. Applied Linear Statistical Models. Regression, Analysis of Variance, and Experimental Designs. Richard D. Irwin Inc. Homewood, Illinois. 841 pp.
- Northeast Oceanographic and Atmospheric Administration and the National Marine Fisheries Service. 1998. Corrigendum. SAW - 26. Advisory Report on Stock Status. Northeast Fisheries Science Center. Woodshole, Massachusetts. pp. 24 - 29.

- Rago, P. J., K. A. Sosenee, J. K. T. Brodziak, S. A. Murawski, and E. D.
   Anderson. 1994. Distribution and dynamics of Atlantic spiny dogfish (*Squalus acanthias*). NOAA/NMFS Ref. Doc. 94-19. 102 pp.
- Ricker, W. E. 1975. J. C. Stevenson (ed.). Computation and Interpretation of Biological Statistics of Fish Populations. Department of the Environment Fisheries and Marine Service, Ottawa.
- Rulifson, R. A. 1998. Biological characterization of the North Carolina spiny dogfish (*Squalus acanthias*) fishery. Interim Report. Fish. Res. Grant Prog. 7 pp. 10.
- Salsbury, J. 1986. Spiny dogfish in Canada. Can. Ind. Rep. Fish. and Aquat.Sci. 169. 57 pp.
- Saunders, M. W. and G. A. McFarlane. 1993. Age and length at maturity of the female spiny dogfish, *Squalus acanthias*, in the Strait of Georgia, British Columbia, Canada. Env. Bio. Fish. 38: 49-57.
- Scott, W. B. and M. G. Scott. 1988. *Atlantic Fishes of Canada*. University of Toronto Press, Toronto.

- Shafer, T. C. 1970. Dogfish migration in the western North Atlantic 1966 1969. Master's thesis. University of Rhode Island.
- Siezen, R. J. 1989. Eye lens aging in the spiny dogfish (*Squalus acanthias*) I. Age determination from lens weight. Cur. Eye Res. 8: 707 - 712.
- Slauson, T. P. 1982. Growth, maturation, and fecundity of the spiny dogfish Squalus acanthias, in the Northwest Atlantic. Master's thesis. State University of New York. Stony Brook, NY. 97 pp.
- Soldat, V. T. 1982. Age and size of spiny dogfish, *Squalus acanthias*, in the northwest Atlantic. Sci. Coun. Stud. 3: 47 52.
- Stearns S. C. and R. E. Crandall. 1984. Plasticity for age and size at sexual maturity: a life-history response to unavoidable stress. pp. 13 - 33. In: G. W. potts and R. J. Wootton (ed.) Fish Reproduction: Strategies and tactics. Academic Press, London.
- Steel, R. G. D. and J. H. Torrie. 1960. Principles and Procedures of StatisticsWith Special Reference to the Biological Sciences. McGraw-Hill BookCompany Ltd., New York. 669 pp.

- Templeman, W. 1944. The life-history of the spiny dogfish (*Squalus acanthias*) and the Vitamin A values of dogfish liver oil. Res. Bull. Dept. Nat. Res, Nfld. 15. 102 pp.
- Templeman, W. 1954. Migrations of spiny dogfish tagged in Newfoundland waters. J. Fish. Res. Bd. Can. 11: 351 -354.
- Templeman, W. 1976. Transatlantic migrations of spiny dogfish (*Squalus acanthias*). J. Fish. Res. Bd. Can. 33: 2605 2609.
- Templeman, W. 1984. Migrations of spiny dogfish, *Squalus acanthias*, and recapture success from tagging in the Newfoundland area, 1963 65. J.
  Northw. Atl. Fish. Sci. 5: 47 53.
- Tuerke, T. M., C. S. Farmer, G. Hamilton, K. Kerle, A. Macy, J. Norby, S.
  Rubenstein, L. Thompson, P. Youtz, R. Forood, and S. Simon. 1993.
  SigmaPlot and SigmaStat Computer Software Program. Jandel Scientific.
  San Rafael, CA.
- Welch, P. S. 1948. Limnological Methods. McGraw-Hill Book Company. New York. 381 pp.

- Wood, C. C., K. S. Ketchen, and R. J. Beamish. 1979. Population dynamics of spiny dogfish (*Squalus acanthias*) in British Columbia waters. J. Fish. Res.
  Bd. Can. 36: 647 656.
- Zigman, S. and T. Yulo. 1979. Eye lens ageing in the dogfish (*Mustelus canis*). Comp. Biocem. Phys. 63B: 379 - 385.

Tables

Dates	Number tagged dogfish	Number sacrificed dogfish	American shad	Winter skate	Striped bass	Method
June 30	<b>Z</b>		10		26	Gillnet
July 7	29	3		5		Longline
July 8			11		11	Gillnet
July 10	81	7		10		Longline
July 12	36	2		2		Longline
July 16	11	1		2		Longline
July 18	85	20		8		Longline
July 19	51	4		7		Longline
July 23	118	9		3		Longline
July 27	68	3		13		Longline
July 28	72	5		6		Longline
Aug. 3	105	12		23		Longline
Aug. 6	73	10		12		Longline
Aug. 8	20	2		2		Longline
Aug. 12		1				Shoreset
Aug. 13		2				Shoreset
Aug. 13		33		2		Longline
Aug. 21		20		4		Longline
Aug. 22	2	39		2		Longline
Aug. 28	244			23		Otter Trawl
Aug. 31		24		2		Longline
Sept. 8		79		8		Longline
Sept. 21		30		4		Longline
Sept. 29		33		4		Longline
Oct. 10		25		2		Longline
TOTAL	995	364	21	144	37	

Table 1: Summary of the fish captured in the Minas Basin during the summer -

-----r

fall 1996.

Table 2:	Summary of total	length range	(cm) and a	ige range (y	years) of	male an	d
	female dogfish ca	ptured in the	three samp	oling locatio	ons.		

	Total length range (cm)		Age range (years)		
	females	males	females	males	
N246 survey	51 – 87	53 – 86	3 – 36	2 – 29	
N249 survey	64 – 101	64 – 87	9 – 27	9 – 24	
Minas Basin	64 - 113	74 – 86	10 – 34	10 – 21	

Table 3: Results of the Kruskal-Wallis one way analysis of variance of ranks between total length (cm) and age (years) of male and female dogfish captured at the three study locations. TL = Total Length

Groups	Median value	Normality	P - value	Significant
				difference
Female TL		····		
N246	71.0	Failed	P < 0.0001	Yes
N249	81.0	P <u>&lt; 0</u> .0001		
Minas Basin	90.0			
Male TL				
N246	71.0	Failed	P < 0.0001	Yes
N249	75.0	P < 0.0001		
Minas Basin	78.5			
Female age				
N24 6	15.0	Failed	P < 0.0001	Yes
N249	18.0	P < 0.0002		
Minas Basin	21.0			
Male age				
N246	15.0	Failed	P = 0.5038	No
N249	16.0	P = 0.0036		
Minas Basin	15.0			

Table 4: Analysis of growth between male and female dogfish in the three study

Comparisons	tcalciculated	Equal growth or not equal growth
FEMALES		
N246 - Minas Basin	0.0982	Equal
N246 - N249	1.0684	Equal
N249 - Minas Basin	1.0563	Equal
MALES		
N246 - Minas Basin	2.2580	Not Equal
N246 - N249	1.0389	Equal
N249 - Minas Basin	1.5774	Equal

locations in Atlantic Canada.

~~~~

|                     | Sex | L      | K    | ťo    |
|---------------------|-----|--------|------|-------|
| Present study       |     |        |      | ····· |
| Minas Basin         | М   | 83     | 0.24 | 2.13  |
|                     | F   | 97     | 0.14 | 2.80  |
| N246 survey         | М   | 77     | 0.27 | 1.98  |
|                     | F   | 80     | 0.20 | 1.90  |
| N249 survey         | М   | 81     | 0.22 | 1.70  |
|                     | F   | 87     | 0.17 | 2.20  |
| All 3 locations     | М   | 78     | 0.25 | 1.80  |
|                     | F   | 91     | 0.17 | 2.50  |
| Slauson 1975        | М   | 85.48  | 0.14 | -1.96 |
|                     | F   | 120.96 | 0.07 | -2.75 |
| Nammack et al. 1985 | М   | 82.49  | 0.18 | -2.67 |
|                     | F   | 100.50 | 0.09 | -2.90 |
|                     |     |        |      |       |

Table 5: The von Bertalanffy growth parameters determined for spiny dogfish for

the present study and other studies in the Northwest Atlantic.
| Percent females | Percent females                                                                 | Percent females                                                                                                  |  |
|-----------------|---------------------------------------------------------------------------------|------------------------------------------------------------------------------------------------------------------|--|
| N246 survey     | N249 survey                                                                     | Minas Basin                                                                                                      |  |
| 95.2            | 72.3                                                                            | 35.8                                                                                                             |  |
| 2.4             | 0                                                                               | 8.1                                                                                                              |  |
| 0               | 16.9                                                                            | 18.4                                                                                                             |  |
| 0               | 1.5                                                                             | 2.9                                                                                                              |  |
| 2.4             | 4.6                                                                             | 24.2                                                                                                             |  |
| 0               | 4.6                                                                             | 10.6                                                                                                             |  |
|                 | Percent females<br>N246 survey<br>95.2<br>2.4<br>0<br>0<br>2.4<br>0<br>2.4<br>0 | Percent females Percent females   N246 survey N249 survey   95.2 72.3   2.4 0   0 16.9   0 1.5   2.4 4.6   0 4.6 |  |

Table 6: Percent number of female dogfish in the reproductive development

stages for the three study locations during the summer - fall 1996.

57

Table 7: Summary of egg diameter ranges (mm), embryo total length range (mm), and male and female dogfish pups total length range (mm) in the various reproductive development stages.

| <br>Reproductive | Egg diameter | Embryo       | Mean embryo  | Pup total length<br>(mm) |          |
|------------------|--------------|--------------|--------------|--------------------------|----------|
| development      | (mm)         | total length | total length |                          |          |
|                  |              | (mm)         | (mm)         | female                   | male     |
| <br>Stage 2      | 20.8 - 41.6  |              | ······       |                          |          |
| Stage 3          | 18.2 - 64.4  | 16.1 - 71.6  | 39.8         |                          |          |
| Stage 4          | 11.4 – 71.6  | 40.4 - 97.8  | 72.3         |                          |          |
| Stage 5          | 17.2 - 43.3  | 176 – 398    | 224.8        | 176 - 326 1              | 74 - 398 |
| Stage 6          | 23.1 - 40.5  |              |              |                          |          |
|                  |              |              |              |                          |          |

Table 8: Summary of tag recoveries from spiny dogfish marked in the Minas

| Tag      | Date      | Date       | Days  | Location      | Distance    |
|----------|-----------|------------|-------|---------------|-------------|
| number   | tagged    | captured   | at    | captured      | travelled   |
|          | (1996)    |            | large |               | <u>(km)</u> |
| 1. 1264  | July 23   | Nov. 2/96  | 93    | 42.10N 70.01W | 801         |
| 2. 1694  | July 19   | Dec. 21/96 | 160   | 43.19N 67.09W | 432         |
| 3. 1190  | July 23   | Dec. 21/96 | 155   | 43.39N 66.57W | 416         |
| 4. 1053  | July 10   | Dec. 1/96  | 144   | 41.06N 71.30W | 1056        |
| 5. 1646  | July 18   | Dec. 29/96 | 164   | 43.00N 65.43w | 554         |
| ** 1264  | July 23   | Nov. 2/96  | 93    | 42.05N 70.15W |             |
| 6. 212   | August 28 | Feb. 25/97 | 153   | 44.05N 66.17W | 1049        |
| 7. 1580  | July 8    | Apr. 9/97  | 275   | 43.50N 65.45W | 395         |
| 8. 1226  | July 27   | May 12/97  | 289   | 42.36N 70.10W | 736         |
| 9. 1425  | July 28   | May 24/97  | 300   | 45.12N 64.20W | 3           |
| 10. 1904 | August 28 | May 3/97   | 248   | 42.37N 70.00W | 736         |
| 11. 1294 | July 23   | Jun. 6/97  | 317   | 43.60N 66.60W | 599         |
| 12.1901  | August 28 | Jul. 12/97 | 318   | 42.40N 71.00W | 820         |
| 13.1480  | July 31   | Jul. 31/97 | 363   | 45.05N 64.10W | 29          |
| 14.1640  | July 18   | Aug. 6/97  | 384   | 44.30N 66.05W | 401         |
| 15. 1274 | July 23   | Aug. 12/97 | 385   | 43.58N 58.66W | 416         |
| 16. 1702 | August 6  | Jun. 10/97 | 308   | 41.54N 68.54W | 881         |
| 17. 1159 | July 23   | May 15/98  | 630   | 44.27N 66.42W | 348         |
| Average  |           |            | 276   |               | 569         |

Basin during July 8 – August 28, 1996.

\*\* Tag captured twice on the same day first by NOAA Survey and returned to the

water, then later by a fisherman who kept the fish and the tag.

Figures

Figure 1: The Minas Basin sampling locations for dogfish captured from July 7, 1996 to October 10, 1996.



•

. •

Figure 2: The N246 DFO survey area and set locations for spiny dogfish captured during July 1996.



.

.

64

Figure 3: The N249 DFO survey area and set locations for dogfish captured during September 1996.



•

.

.

66

:

Figure 4: Posterior dorsal spine removal in the spiny dogfish for an age estimate.

a = first cut; b = second cut.



•

:

Figure 5: Longitudinal schematic diagram of the posterior dorsal spine (a) and spine growth showing five annual opaque rings or bands at the cone bases (b)(Slauson 1982).



.

Figure 6: Measurements taken from the posterior dorsal spine of the spiny dogfish. SL = Spine Length; WL = Wear Length; SBD = Spine Base Diameter; and WPD = Wear Point Diameter .



Figure 7: Reproductive tract removal in the female spiny dogfish.



.

Figure 8: Stage I of reproductive tract development in the female spiny dogfish.

•



.

Figure 9: Stage II of reproductive tract development in the female spiny dogfish.



t

Figure 10: Stage III of reproductive tract development in the female spiny dogfish.



ŧ

Figure 11: Stage IV of reproductive tract development in the female spiny

dogfish.



-

.

Figure 12: Stage V of reproductive tract development in the female spiny

dogfish.



-

\$

Figure 13: Stage VI of reproductive tract development in the female spiny

dogfish.



۰.

Figure 14: The N246 DFO Scotian Shelf survey set locations and number of dogfish captured in each tow during July 1996.



•

· ·

٠.

٩

•

88

Figure 15: The N249 DFO Gulf of St. Lawrence survey set locations and number

of spiny dogfish captured in each tow during September 1996.



. .

-

.

.

90

.

Figure 16: Total length frequency of female dogfish captured in the Minas Basin, during the N246 DFO survey, and during the N249 DFO survey, summer - fall 1996.


Figure 17: Total length frequency of male dogfish captured in the Minas Basin, during the N246 DFO survey, and during the N249 DFO survey, summer - fall 1996.



Figure 18: Age frequency of female dogfish captured in the Minas Basin, during the N246 DFO survey, and during the N249 DFO survey, summer - fall 1996.



Figure 19: Age frequency of male dogfish captured in the Minas Basin, during the N246 DFO survey, and during the N249 DFO suvey, summer - fall 1996.



Figure 20: Weight-length relationship of female spiny dogfish captured in the Minas Basin, during the N246 DFO survey, and during the N249 DFO survey, summer-fall 1996.



Figure 21: Weight-length relationship of male spiny dogfish captured in the Minas Basin, during the N246 DFO survey, and during the N249 DFO survey, summer- fall 1996.



Figure 22: Weight-length relationship of male and female spiny dogfish caught at the three study locations in Atlantic Canada during the summer fall 1996.



Figure 23: Spine base diameter (mm) in relation to age (years) of male and female dogfish with wear point diameters ≤1.5 mm to determine the equation for lost annuli due to wear.

-



Figure 24: The von Bertalanffy growth equations for female dogfish captured in the Minas Basin, during the N246 DFO survey, and during the N249 DFO survey, summer - fall 1996. Bars represent 1 standard deviation from the mean.

.



Figure 25: The von Bertalanffy growth equations for male dogfish captured in the Minas Basin, during the N246 DFO survey, and during the N249 DFO survey, summer - fall 1996. Bars represent 1 standard deviation from the mean.



Figure 26: The von Bertalanffy growth equations for male and female dogfish captured at the three study locations in Atlantic Canada during the summer - fall 1996. Bars represent 1 standard deviation from the mean.



Figure 27: Percent maturity of female dogfish captured in the Minas Basin, during the N246 DFO survey, and during the N249 DFO survey, summer - fall 1996.



Figure 28: Percent maturity of female dogfish captured at the three study

locations in Atlantic Canada during the summer - fall 1996.



Figure 29: Number of progeny (NP) in relation to total maternal weight(TMW)(kg) for fecundity of the female dogfish captured in the Minas Basin, on the N246 DFO survey, and on the N249 DFO survey during the summer - fall 1996.



Figure 30: Number of progeny (NP) in relation to the total maternal weight (TMW)(kg) for fecundity of female dogfish captured at the three study locations in Atlantic Canada during the summer - fall 1996.



Figure 31: Tag return locations for dogfish marked in the Minas Basin during the summer of 1996.

•



Appendices

## Appendix 1: Raw data for female spiny dogfish captured in the Minas Basin from July 7 - October 10, 1996. TL = Total

| Label | TL<br>(cm) | FL      | Weight | Counted | Calculated | Reproductive | Egg    | Embryo |
|-------|------------|---------|--------|---------|------------|--------------|--------|--------|
|       |            | m) (cm) | (kg)   | age     | age        | stage        | number | number |
| 1b    | 100        | 90      | 4.3    | 10      | 22         | 3            | -      | 8      |
| 2b    | 103        | 91      | 4.5    | 9       | 22         | 3            | -      | 11     |
| 3b    | 84         | 85      | 3.6    | 10      | 25         | 5            | 7      | 6      |
| 5b    | 95         | 86      | 4.0    | 16      | 19         | 5            | 7      | 6      |
| 6b    | 94         | 86      | 3.5    | 13      | 24         | 5            | 4      | 1      |
| 7b    | 97         | 85      | 4.1    | 9       | 21         | 5            | 7      | 5      |
| 8b    | 102        | 89      | 3.9    | 10      | 24         | 5            | 6      | 2      |
| 9b    | 91         | 81      | 3.5    | 13      | 20         | -            | -      | -      |
| 10b   | 85         | 75      | 2.5    | 14      | -          | 1            | -      | -      |
| 11b   | 77         | 67      | 1.8    | 15      | 17         | 1            | -      | -      |
| 12b   | 97         | 86      | 4.2    | 14      | 23         | 5            | 7      | 5      |
| 13b   | 84         | 74      | 3.0    | 18      | 27         | 3            | -      | 5      |
| 14b   | 102        | 92      | 5.0    | 8       | 22         | 3            | -      | 2      |
| 15b   | 100        | 89      | 5.0    | 12      | 21         | -            | -      | -      |
| 16b   | 97         | 86      | 3.6    | 10      | 21         | 3            | -      | 5      |
| 17b   | 96         | 85      | 3.8    | 14      | 22         | -            | -      | -      |
| 18b   | 94         | 84      | 3.9    | 18      | 27         | 3            | -      | 6      |
| 19b   | 94         | 83      | 3.7    | 17      | 24         | 5            | 4      | 5      |
| 20b   | 88         | 77      | 3.3    | 15      | 21         | 3            | -      | 4      |
| 21b   | 98         | 83      | 3.9    | 7       | 21         | 3            | -      | 3      |
| 22b   | 92         | 80      | 3.9    | 13      | 20         | 5            | 6      | 6      |
| 23b   | 91         | 80      | 3.4    | 12      | 19         | 5            | 5      | 6      |
| 24b   | 87         | 76      | 3.2    | 11      | 16         | 2            | 5      | -      |
| 25b   | 93         | 82      | 3.7    | 18      | 24         | 3            | -      | 5      |

length; FL = Fork length.

124

| Label | TL   | FL             | Weight | Counted | Calculated | Reproductive | Egg    | Embryo |
|-------|------|----------------|--------|---------|------------|--------------|--------|--------|
|       | (cm) | (cm)           | (kg)   | age     | age        | stage        | number | number |
| 26b   | 99   | 87             | 4.6    | 11      | 21         | 3            | -      | 5      |
| 27b   | 94   | 82             | 3.4    | 13      | 23         | 3            | -      | 3      |
| 28b   | 94   | 83             | 4.0    | 14      | 24         | 3            | -      | 4      |
| 29b   | 86   | 76             | 2.7    | 17      | 22         | 3            | -      | 2      |
| 30b   | 89   | 79             | 3.3    | 16      | 23         | 3            | -      | 2      |
| 31b   | 93   | 81             | 3.7    | 16      | 20         | 3            | -      | 4      |
| 32b   | 105  | 94             | 5.2    |         |            | 3            | -      | 6      |
| 33b   | 90   | 7 <del>9</del> | 3.0    | 6       | 18         | 5            | 5      | 5      |
| 34b   | 95   | 83             | 3.7    | 12      | 18         | 3            | -      | 4      |
| 35b   | 98   | 88             | 5.5    | 10      |            | 5            | 7      | 7      |
| 36b   | 95   | 85             | 4.5    | 14      | 23         | 3            | -      | 6      |
| 37b   | 79   | 68             | 1.8    | 26      | 29         | 1            | -      | -      |
| 38b   | 73   | 64             | 1.7    | 17      | 17         | 1            | -      | -      |
| 39b   | 89   | 74             | 2.8    | 17      | 21         | 2            | 5      | -      |
| 40b   | 86   | 75             | 3.1    | 30      | 33         | 3            | -      | 2      |
| 41b   | 94   | 84             | 3.5    | 14      | 22         | -            | -      | -      |
| 42b   | 95   | 83             | 4.0    | 12      | 22         | 3            | -      | 3      |
| 43b   | 97   | 85             | 4.6    | 18      | 24         | 5            | 7      | 7      |
| 44b   | 88   | 78             | 3.4    | 15      | 21         | 3            | -      | 4      |
| 45b   | 86   | 76             | 3.2    | 18      | 23         | 3            | -      | 5      |
| 46b   | 102  | 90             | 4.6    | 19      | 29         | 3            | -      | 6      |
| 47b   | 98   | 87             | 3.9    | 15      | 23         | 3            | -      | 5      |
| 48b   | 100  | 90             | 5.0    | 6       | 23         | -            | -      | -      |
| 49b   | 89   | 79             | 3.4    | 17      | 25         | 5            | 6      | 3      |
| 50b   | 97   | 86             | 4.3    | 21      | 27         | 2            | 3      | -      |

125

| Label | TL<br>(cm) | FL             | . Weight<br>1) (kg) | Counted Calculated<br>age age | Reproductive | Ega   | Embryo |        |
|-------|------------|----------------|---------------------|-------------------------------|--------------|-------|--------|--------|
|       |            | (cm)           |                     |                               | age          | stage | number | number |
| 51b   | 93         | 81             | 4.3                 | 13                            | 24           |       |        | -      |
| 52b   | 102        | 90             | 4.7                 | 12                            | 21           | 5     | 8      | 3      |
| 53b   | 104        | 91             | 5.2                 | 10                            | 23           | -     | -      | -      |
| 54b   | 100        | 87             | 4.4                 | 8                             | 21           | -     | -      | -      |
| 55b   | 83         | 73             | 2.5                 | 18                            | 20           | -     | -      | -      |
| 56b   | 109        | 97             | 5.8                 | 13                            | 19           | 3     | -      | 4      |
| 57b   | 93         | 82             | 4.0                 | 20                            | 28           | 5     | 7      | 3      |
| 58b   | 85         | 74             | 2.7                 | 21                            | 26           | 3     | -      | 3      |
| 59b   | 83         | 72             | 2.5                 | 28                            | 30           | 1     | -      | -      |
| 60b   | 87         | 77             | 2.9                 | 19                            | 24           | 1     | -      | -      |
| 61b   | 96         | 85             | 4.3                 |                               |              | 3     | -      | 5      |
| 62b   | 81         | 72             | 2.6                 | 12                            | 16           | -     | -      | -      |
| 63b   | 99         | 88             | 4.5                 | 25                            | 31           | 3     | -      | 4      |
| 64b   | 103        | 91             | 5.2                 | 11                            | 24           | 3     | -      | 4      |
| 65b   | 80         | 70             | 2.5                 | 22                            | 25           | 2     | 6      | -      |
| 66b   | 90         | 7 <del>9</del> | 3.4                 | 15                            | 20           | 5     | 5      | 4      |
| 67b   | 99         | 87             | 4.5                 | 11                            | 26           | 3     | -      | 4      |
| 68b   | 83         | 73             | 2.4                 | 19                            | 23           | 3     | -      | 8      |
| 69b   | 83         | 73             | 2.8                 | 29                            | 34           | 3     | -      | 8      |
| 70b   | 103        | 91             | 5.8                 | 8                             | 19           | 3     | -      | 8      |
| 71b   | 100        | 87             | 4.7                 | 16                            | 28           | 3     | -      | 8      |
| 72b   | 92         | 81             | 3.8                 | 10                            | 23           | 5     | 7      | 4      |
| 73b   | 101        | 87             | 4.6                 | 14                            | 24           | 3     | -      | 4      |
| 74b   | 97         | 87             | 4.1                 | 20                            | 28           | 3     | -      | 4      |
| 75b   | 94         | 83             | 4.3                 | 17                            | 24           | 5     | 7      | 7      |

| Label | TL<br>(cm)     | FL<br>(cm) | Weight      | Counted | Calculated | Reproductive | Egg    | Embryo |
|-------|----------------|------------|-------------|---------|------------|--------------|--------|--------|
| 765   |                | 77         | <u>(ry/</u> | aye     | aye        | Slaye        | number | number |
| 700   | 00             | 77         | 3.1         | 10      | 24         | 3            | -      | 3      |
| //D   | 84             | /5         | 2.7         | 14      | 19         | 3            | -      | 4      |
| 78b   | 95             | 84         | 4.2         | 15      | 23         | -            | -      | -      |
| 79b   | 100            | 89         | 4.0         | 11      | 22         | 3            | -      | 1      |
| 80b   | <del>9</del> 7 | 86         | 5.1         | 16      | 21         | -            | -      | -      |
| 81b   | 90             | 80         | 3.4         | 23      | 27         | 3            | -      | 5      |
| 82b   | 83             | 73         | 2.7         | 15      | 19         | 2            | 2      | -      |
| 83b   | 95             | 83         | 3.7         | 17      | 22         | 6            | 5      | -      |
| 84b   | 85             | 74         | 3.2         | 15      | 24         | 5            | 5      | 5      |
| 85b   | 81             | 72         | 2.6         | 16      | 20         | 1            | -      | 4      |
| 86b   | 80             | 72         | 2.8         | 16      | 22         | 2            | 5      | -      |
| 87b   | 90             | 80         | 3.7         | 20      | 28         | 3            | -      | 4      |
| 88b   | 77             | 68         | 1.9         | 19      | 19         | 3            | -      | Å      |
| 89b   | 71             | 62         | 1.5         | 15      | 15         | 1            | -      | -      |
| 90b   | 90             | 80         | 3.4         |         |            | 5            | 4      | 4      |
| 91b   | 79             | 69         | 2.2         | 22      | 25         | 2            | 4      | -      |
| 92b   | 86             | 76         | 2.7         | 26      | 28         | 2            | 6      | -      |
| 93b   | 87             | 76         | 3.6         | 23      | 27         | 5            | 7      | 5      |
| 94b   | 86             | 77         | 3.3         |         |            | 5            | 5      | 4      |
| 95b   | 95             | 85         | 4.6         | 16      | 27         | 5            | 7      | 5      |
| 96b   | 84             | 75         | 28          | 21      | 26         | 2            | ,<br>6 | 11     |
| 97h   | 93             | 83         | 3.5         | 20      | 25         | 5            | 5      | 2      |
| QRh   | 94             | 84         | 3.0         | 27      | LU         | 5            | 7      | 2      |
| 00b   | 78             | 68         | 1.8         | 18      | 21         | 5            | 1      | 3      |
| 100h  | 24             | 70         | 1.0         | 47      | 21         | 3<br>4       | -      | 4      |
|       | 01             | / V        | ۷.۷         | I /     |            | 1            |        | -      |

127

| Label | TL   | FL   | Weight | Counted | Calculated | Reproductive | Egg    | Embryo |
|-------|------|------|--------|---------|------------|--------------|--------|--------|
|       | (cm) | (cm) | (kg)   | age     | age        | stage        | number | number |
| 101b  | 94   | 83   | 5.2    | 13      | 23         | 5            | 6      | 7      |
| 102b  | 93   | 82   | 4.4    | 14      | 25         | 5            | 7      | 7      |
| 103b  | 90   | 80   | 4.0    | 17      | 23         | 5            | 7      | 6      |
| 104b  | 88   | 78   | 3.1    | 15      | 21         | 3            | -      | 6      |
| 105b  | 85   | 75   | 2.8    | 14      | 20         | 6            | 5      | -      |
| 106b  | 83   | 73   | 2.7    | 15      | 18         | 2            | 4      | -      |
| 107b  | 90   | 79   | 3.7    | 18      | 22         | 2            | 6      | -      |
| 108b  | 92   | 82   | 4.5    | 11      | 20         | 5            | 8      | 6      |
| 109b  | 90   | 80   | 3.9    | 10      | 19         | 5            | 7      | 6      |
| 110b  | 82   | 72   | 2.8    | 13      | 17         | 6            | -      | 5      |
| 111b  | 84   | 75   | 2.7    | 12      | 20         | 3            | -      | 5      |
| 112b  | 85   | 75   | 2.7    | 14      | 18         | 3            | -      | -      |
| 113b  | 84   | 75   | 2.8    | 10      | 14         | 6            | -      | 5      |
| 114b  | 73   | 64   | 1.5    | 10      | 10         | 1            | -      | -      |
| 115b  | 86   | 77   | 3.2    | 16      | 23         | 3            | -      | 4      |
| 116b  | 95   | 84   | 4.2    | 10      | 22         | 5            | 5      | 2      |
| 117b  | 86   | 76   | 2.9    | 11      | 17         | 1            | -      | -      |
| 118b  | 92   | 82   | 4.2    | 19      | 23         | 5            | 8      | 4      |
| 119b  | 77   | 68   | 1.8    | 10      | 22         | 1            | -      | -      |
| 120b  | 81   | 72   | 2.2    | 17      | 23         | -            | -      | -      |
| 121b  | 78   | 69   | 2.0    | 12      | 12         | 1            | -      | -      |
| 122b  | 83   | 73   | 2.7    | 17      | 20         | 6            | 4      | -      |
| 123b  | 87   | 77   | 3.1    | 16      | 20         | -            | -      | -      |
| 124b  | 87   | 76   | 3.0    | 19      | 25         | 1            | -      | -      |
| 125b  | 80   | 71   | 2.1    | -       | •          | 1            | -      | -      |

128

----
| Label | TL   | FL   | Weight | Counted | Calculated | Reproductive | Eaa    | Embrvo |
|-------|------|------|--------|---------|------------|--------------|--------|--------|
|       | (cm) | (cm) | (kg)   | age     | age        | stage        | number | number |
| 127b  | 75   | 67   | 1.8    | 14      | 16         | 1            | -      | -      |
| 128b  | 77   | 67   | 2.0    | 21      | 21         | 1            | -      | -      |
| 129b  | 80   | 71   | 2.1    | 15      | 19         | 1            | -      | -      |
| 130b  | 94   | 82   | 3.7    | 10      | 21         | 5            | 5      | 5      |
| 131b  | 87   | 77   | 3.3    | 15      | 24         | 5            | 5      | 5      |
| 132b  | 76   | 67   | 2.0    | 14      | 15         | 1            | -      | -      |
| 133b  | 84   | 74   | 2.7    | 16      | 19         | 1            | -      | -      |
| 134b  | 81   | 71   | 2.3    | 18      | 21         | 1            | -      | -      |
| 135b  | 75   | 66   | 2.1    | 17      | 17         | 1            | -      | -      |
| 136b  | 72   | 63   | 1.6    | 18      | 19         | 1            | -      | -      |
| 137b  | 80   | 70   | 2.5    | 14      | 19         | 1            | -      | -      |
| 138b  | 74   | 66   | 1.7    | 13      | 13         | 1            | -      | -      |
| 139b  | 79   | 70   | 2.0    | 18      | 20         | 1            | -      | -      |
| 140b  | 102  | 90   | 5.0    | 14      | 15         | 3            | -      | 6      |
| 141b  | 85   | 75   | 3.0    | 14      | 21         | 1            | -      | -      |
| 142b  | 81   | 72   | 2.6    | 19      | 22         | 2            | 4      | -      |
| 143b  | 73   | 64   | 1.7    | 19      | 21         | 1            | -      | -      |
| 144b  | 80   | 71   | 2.0    | 15      | 16         | 1            | -      | -      |
| 145b  | 75   | 65   | 1.9    | 16      | 18         | 1            | -      | -      |
| 146b  | 80   | 70   | 2.1    | 18      | 19         | -            | -      | -      |
| 147b  | 82   | 71   | 2.9    | 14      | 16         | 2            | 5      | -      |
| 148b  | 95   | 83   | 4.0    | 13      | 23         | 3            | -      | 2      |
| 149b  | 91   | 80   | 3.5    | 17      | 22         | 2            | 7      | -      |
| 150b  | 81   | 72   | 2.2    | 16      | 20         | 1            | -      | -      |
| 151b  | 99   | 90   | 5.1    | 13      | 25         | 1            | 7      | 9      |

| Label | TL   | FL             | Weight | Counted | Calculated | Reproductive | Egg    | Embryo |
|-------|------|----------------|--------|---------|------------|--------------|--------|--------|
|       | (cm) | (cm)           | (kg)   | age     | age        | stage        | number | number |
| 152b  | 77   | 67             | 1.9    | 11      | 17         | 1            | -      | -      |
| 153b  | 86   | 76             | 2.6    | 12      | 19         | 1            | -      | -      |
| 154b  | 80   | 72             | 2.5    | 17      | 18         | 1            | -      | -      |
| 155b  | 79   | 69             | 1.9    | 15      | 18         | 1            | -      | -      |
| 156b  | 86   | 77             | 3.3    | 19      | 24         | 2            | 2      | -      |
| 157b  | 85   | 76             | 3.0    | 16      | 19         | -            | -      | -      |
| 158b  | 67   | 59             | 1.4    | 15      | 15         | 1            | -      | -      |
| 159b  | 73   | 63             | 1.5    | 15      | 15         | 1            | -      | -      |
| 160b  | 82   | 75             | 2.1    | 15      | 24         | 2            | 3      | -      |
| 161b  | 95   | 85             | 4.3    | 14      | 20         | 5            | 7      | 6      |
| 162b  | 87   | 77             | 3.4    | 14      | 21         | 3            | -      | 5      |
| 163b  | 90   | 81             | 3.5    | 17      | 22         | 2            | 3      | -      |
| 164b  | 78   | 6 <del>9</del> | 2.0    | 15      | 18         | 1            | -      | -      |
| 165b  | 70   | 62             | 1.4    | 12      | 14         | 1            | -      | -      |
| 166b  | 81   | 71             | 2.3    | 10      | 14         | 1            | -      | -      |
| 167b  | 82   | 71             | 2.5    | 15      | 16         | 1            | -      | -      |
| 168b  | 95   | 84             | 3.5    | 15      | 21         | 3            | -      | 9      |
| 169b  | 78   | 69             | 2.1    | 21      | 23         | 1            | -      | •      |
| 170b  | 83   | 73             | 3.9    | 11      | 19         | 5            | 5      | 5      |
| 171b  | 80   | 71             | 2.0    | 18      | 19         | 1            | -      | -      |
| 172b  | 75   | 66             | 1.7    | 14      | 17         | 1            | -      | -      |
| 173b  | 79   | 70             | 2.0    | 14      | 17         | 1            | -      | -      |
| 174b  | 72   | 63             | 1.5    |         |            | 1            | -      | -      |
| 175b  | 89   | 79             | 3.4    | 12      | 19         | -            | -      | -      |
| 176b  | 81   | 71             | 2.5    | 14      | 16         | 1            | -      | -      |

130

....

-

| Label | TL   | FL   | Weight | Counted | Calculated | Reproductive | Eaa    | Embryo |
|-------|------|------|--------|---------|------------|--------------|--------|--------|
|       | (cm) | (cm) | (kg)   | age     | age        | stage        | number | number |
| 177b  | 84   | 74   | 3.2    | 19      | 23         | 2            | 4      | -      |
| 178b  | 82   | 72   | 2.4    | 18      | 20         | -            | -      | -      |
| 179b  | 87   | 77   | 3.3    | 17      | 21         | 2            | 4      | -      |
| 180b  | 88   | 77   | 2.8    | 15      | 19         | 2            | 4      | -      |
| 181b  | 104  | 93   | 4.9    | 6       | 21         | 3            | -      | 7      |
| 182b  | 79   | 70   | 2.4    | 18      | 19         | 1            | -      | -      |
| 183b  | 100  | 89   | 5.1    | 13      | 19         | -            | -      | -      |
| 184b  | 93   | 82   | 4.3    | 16      | 21         | 5            | 7      | 6      |
| 185b  | 80   | 70   | 2.8    | 18      | 21         | 2            | 4      | -      |
| 186b  | 74   | 64   | 1.8    | 13      | 17         | 1            | -      | -      |
| 187b  | 88   | 77   | 3.3    | 11      | 19         | 5            | 3      | 3      |
| 188b  | 88   | 78   | 2.1    | 13      | 22         | 1            | -      | -      |
| 189b  | 85   | 75   | 3.1    | 20      | 26         | 5            | 3      | 6      |
| 190b  | 87   | 76   | 3.8    | 22      | 26         | 2            | 5      | -      |
| 191b  | 83   | 74   | 3.1    | 15      | 20         | 2            | 5      | -      |
| 192b  | 111  | 100  | 8.0    | -       | -          | 5            | 11     | 8      |
| 193b  | 90   | 79   | 3.2    | 12      | 20         | 5            | 5      | 3      |
| 194b  | 81   | 72   | 2.2    | -       | -          | 1            | -      | -      |
| 195b  | 80   | 70   | 2.4    | 22      | 26         | 1            | -      | -      |
| 196b  | 106  | 94   | 5.7    | 13      | 25         |              | -      | -      |
| 197b  | 102  | 92   | 5.4    | 9       | 20         | 3            | 8      | -      |
| 198b  | 97   | 86   | 4.6    | 14      | 23         | 5            | 6      | 6      |
| 199b  | 78   | 68   | 2.5    | 12      | 17         | •            | -      | -      |
| 200b  | 84   | 73   | 3.4    | 13      | 17         | 6            | 5      | -      |
| 201b  | 90   | 78   | 4.2    | 13      | 23         | 5            | 7      | 5      |

| Label | TL   | FL   | Weight | Counted | Calculated | Reproductive | Egg    | Embryo |
|-------|------|------|--------|---------|------------|--------------|--------|--------|
|       | (cm) | (cm) | (kg)   | age     | age        | stage        | number | number |
| 202b  | 94   | 82   | 4.3    | 10      | 20         | 5            | 17     | 16     |
| 203b  | 81   | 71   | 2.7    | 14      | 18         | 1            | -      | -      |
| 204b  | 100  | 86   | 5.2    | 14      | 24         | 5            | 6      | 5      |
| 205b  | 85   | 73   | 2.8    | 18      | 21         | 1            | -      | -      |
| 206b  | 83   | 73   | 2.7    | 15      | 21         | 6            | 4      | -      |
| 207b  | 86   | 75   | 3.0    | 19      | 23         | 1            | -      | -      |
| 208b  | 82   | 71   | 2.5    | 16      | 20         | 1            | -      | -      |
| 209b  | 91   | 79   | 3.2    | 15      | 21         |              | -      | -      |
| 210b  | 87   | 76   | 2.8    | 21      | 26         | 3            | -      | 5      |
| 211b  | 91   | 80   | 3.5    | 15      | 21         | 6            | 4      | -      |
| 212b  | 68   | 60   | 1.5    | 12      | 12         | 1            | -      | -      |
| 213b  | 74   | 64   | 2.0    | 14      | 16         | 1            | -      | -      |
| 214b  | 88   | 77   | 3.0    | 19      | 23         | 6            | 4      | -      |
| 215b  | 75   | 65   | 2.2    | 17      | 18         | 1            | -      | -      |
| 216b  | 98   | 87   | 5.2    | 10      | 19         | 5            | 7      | 8      |
| 217b  | 80   | 69   | 2.6    | 16      | 20         | 1            | -      | -      |
| 218b  | 78   | 69   | 2.4    |         |            | 6            | 3      | -      |
| 219b  | 72   | 63   | 1.6    | 14      | 15         |              | -      | -      |
| 220b  | 94   | 81   | 4.8    | 17      | 24         | 5            | 6      | 6      |
| 221b  | 77   | 67   | 2.0    | 19      | 20         | 1            | -      | -      |
| 222b  | 88   | 77   | 3.7    | 18      | 23         | 6            | 6      | -      |
| 223b  | 83   | 73   | 3.1    | 15      | 20         | 6            | 5      | -      |
| 224b  | 85   | 75   | 2.9    |         |            | 1            | -      | -      |
| 225b  | 90   | 79   | 3.8    | 11      | 19         | 5            | 6      | 4      |
| 226b  | 83   | 73   | 2.8    | 17      | 22         | 6            | 4      | -      |

| Label | TL<br>(cm) | FL<br>(cm) | Weight      | Counted | Calculated | Reproductive | Egg    | Embryo |
|-------|------------|------------|-------------|---------|------------|--------------|--------|--------|
| 0075  |            |            | <u>(ky)</u> | aye     | aye        | stage        | number | number |
| 227D  | 79         | 70         | 2.3         | 20      | 25         | 1            | -      | -      |
| 228b  | 86         | 76         | 3.0         | 17      | 19         | 6            | 5      | -      |
| 229b  | 71         | 62         | 1.5         | 12      | 12         |              | -      | -      |
| 230b  | 86         | 76         | 3.3         | 16      | 20         | 6            | 5      | -      |
| 231b  | 90         | 80         | 3.7         | 15      | 20         | 6            | 5      | -      |
| 232b  | 69         | 60         | 1.4         | 13      | 12         | 1            | -      | -      |
| 233b  | 87         | 77         | 3.0         | 19      | 23         | 1            | -      | -      |
| 234b  | 84         | 75         | 3.0         | 14      | 18         | 3            | -      | 3      |
| 235b  | 83         | 73         | 2.9         | 16      | 22         | 6            | 6      | -      |
| 236b  | 89         | 78         | 3.4         | 16      | 22         | 5            | 8      | 5      |
| 237b  | 69         | 60         | 1.5         | 14      | 14         | 1            | -      | -      |
| 238b  | 92         | 81         | 4.4         | 11      | 20         | 5            | 5      | 8      |
| 239b  | 75         | 65         | 1.8         | 13      | 14         | -            | -      | -      |
| 240b  | 72         | 63         | 1.7         | 15      | 16         | 1            | -      | -      |
| 241b  | 82         | 72         | 2.7         | 13      | 19         | 1            | -      | -      |
| 242b  | 77         | 68         | 2.1         | 14      | 17         | 1            | -      | -      |
| 243b  | 73         | 64         | 1.6         | 18      | 19         | 1            | -      | -      |
| 244b  | 81         | 70         | 2.4         | 15      | 19         | 6            | 4      | -      |
| 245b  | 80         | 71         | 2.2         | 17      | 18         | 1            | -      | -      |
| 246b  | 73         | 64         | 1.5         | 15      | 16         | 1            | -      | -      |
| 247b  | 84         | 74         | 2.8         | 15      | 19         | 1            | -      | -      |
| 248b  | 81         | 72         | 2.6         | 16      | 19         | 1            |        | _      |
| 249h  | 83         | 74         | 27          | 21      | 25         | ĥ            | ٨      | -      |
| 2506  | 78         | 69         | 21          | 47      | 10         | 4            | 4      | -      |
| 2500  | 10         | 75         | 2.1         | 17      | 10         |              | -      | -      |
| 2310  | 00         | [ ]        | J.U         |         | <u> </u>   | 0            | 5      | -      |

| Label | TL   | FL   | Weight | Counted | Calculated | Reproductive | Egg    | Embryo |
|-------|------|------|--------|---------|------------|--------------|--------|--------|
|       | (cm) | (cm) | (kg)   | age     | age        | stage        | number | number |
| 252b  | 96   | 84   | 4.6    | 9       | 24         | 5            | 6      | 9      |
| 253b  | 87   | 77   | 3.2    | 20      | 25         | 6            | 5      | -      |
| 254b  | 82   | 73   | 2.6    | 15      | 21         | -            | -      | -      |
| 255b  | 83   | 73   | 3.6    | 20      | 22         | 5            | 4      | 2      |
| 256b  | 85   | 75   | 2.8    | 15      | 17         | 1            | -      | -      |
| 257b  | 90   | 79   | 2.9    | -       | -          | 5            | 6      | 5      |
| 258b  | 82   | 74   | 2.5    | 20      | 21         | 1            | -      | -      |
| 259b  | 102  | 90   | 6.5    | 13      | 23         | -            | -      | -      |
| 260b  | 76   | 66   | 1.8    | 16      | 17         | 1            | -      | -      |
| 261b  | 90   | 79   | 4.4    | 17      | 23         | 5            | 6      | 6      |
| 262b  | 89   | 79   | 3.3    | 19      | 23         | 6            | 5      | -      |
| 263b  | 82   | 72   | 2.7    | 15      | 16         | 1            | -      | -      |
| 264b  | 86   | 78   | 3.1    | 12      | 15         | 1            | -      | -      |
| 265b  | 94   | 84   | 4.7    | 24      | 31         | 5            | 7      | 2      |
| 266b  | 80   | 71   | 2.4    | 19      | 21         | 6            | 4      | -      |
| 267b  | 80   | 71   | 2.5    | 14      | 15         | 6            | 4      | -      |
| 268b  | 87   | 77   | 3.1    | 17      | 20         | 1            | -      | -      |
| 269b  | 78   | 69   | 2.4    | 15      | 17         | 1            | -      | -      |
| 270b  | 88   | 79   | 3.4    | 13      | 22         | 1            | -      | -      |
| 271b  | 83   | 74   | 2.4    | 15      | 21         | 1            | -      | -      |
| 272b  | 90   | 79   | 3.0    | 25      | 30         | 1            | -      | -      |
| 273b  | 75   | 66   | 1.7    | 16      | 19         | 1            | -      | -      |
| 274b  | 81   | 71   | 2.5    | 17      | 20         | 6            | 3      | -      |
| 275b  | 74   | 66   | 1.7    | 15      | 18         | 1            | -      | -      |
| 276b  | 79   | 70   | 2.3    | 16      | 19         | 1            | -      | -      |
| 277b  | 73   | 64   | 1.7    | -       | -          | 1            | -      | -      |

| Label        | TL             | FL             | Weight | Counted | Calculated | Reproductive | Egg    | Embryo |
|--------------|----------------|----------------|--------|---------|------------|--------------|--------|--------|
|              | (cm)           | (cm)           | (kg)   | age     | age        | stage        | number | number |
| 278b         | 88             | 79             | 3.1    | 23      | 29         | 4            | 6      |        |
| 279b         | 78             | 69             | 2.6    | 15      | 19         | 1            | -      | -      |
| 280b         | 96             | 86             | 4.4    | 14      | 21         | 5            | 6      | 4      |
| 281b         | 89             | 79             | 3.4    | 17      | 23         | 6            | 6      | -      |
| 282b         | 96             | 86             | 4.5    | 13      | 20         | 5            | 8      | 6      |
| 283b         | 76             | 69             | 2.2    | 19      | 20         | 1            | -      | -      |
| 284b         | 97             | 87             | 5.9    | 10      | 27         | 5            | 7      | 7      |
| 285b         | 83             | 7 <del>9</del> | 3.4    | 16      | 22         | 6            | 5      | -      |
| 286b         | 7 <del>9</del> | 71             | 2.4    | 13      | 19         | 1            | -      | -      |
| 287b         | 85             | 78             | 3.3    | 16      | 22         | 1            | -      | -      |
| 288b         | 83             | 74             | 3.0    | 12      | 18         | 6            | 5      | -      |
| 289b         | 76             | 67             | 1.9    | 16      | 20         | 1            | -      | -      |
| 290b         | 75             | 66             | 2.0    | 13      | 14         | 1            | -      | -      |
| 291b         | 71             | 63             | 1.5    | 13      | 15         | -            | -      | -      |
| 292b         | 91             | 82             | 4.2    | 15      | 25         | 5            | 8      | 5      |
| 293b         | 69             | 60             | 1.8    | 12      | 11         | 1            | -      | -      |
| 294b         | 84             | 76             | 3.6    | 19      | 20         | 6            | 7      | -      |
| 295b         | 85             | 76             | 3.1    | 18      | 28         | 6            | 5      | -      |
| 296b         | 100            | 90             | 6.8    | 11      | 25         | 5            | 9      | 9      |
| <b>29</b> 7b | 100            | 89             | 5.5    | 14      | 20         | 6            | 9      | 9      |
| 298b         | 87             | 77             | 4.2    | 12      | 24         | 5            | 7      | 4      |
| 299b         | 84             | 74             | 3.1    | 21      | 22         | 6            | 5      | -      |
| 300b         | 69             | 63             | 1.6    | 10      | 11         | 1            | -      | -      |
| 301b         | 82             | 74             | 2.8    | 19      | 20         | -            | -      | -      |
| 302b         | 76             | 68             | 2.1    | 18      | 19         | -            | -      | -      |

| Labei | TL   | FL   | Weight | Counted | Calculated | Reproductive | Egg    | Embryo |
|-------|------|------|--------|---------|------------|--------------|--------|--------|
|       | (cm) | (cm) | (kg)   | age     | age        | stage        | number | number |
| 303b  | 100  | 90   | 4.6    | •       | -          | 4            | 6      | 7      |
| 304b  | 77   | 66   | 2.1    | 16      | 18         | 1            | -      | -      |
| 305b  | 68   | 62   | 2.0    | 14      | 19         | 1            | -      | -      |
| 306b  | 100  | 89   | 5.5    | 18      | 28         | 6            | -      | 4      |
| 307b  | 104  | 95   | 5.5    | -       | -          | 5            | 8      | 8      |
| 308b  | 101  | 89   | 4.9    | 12      | 25         | 5            | 6      | 5      |
| 309b  | 71   | 64   | 1.6    | 15      | 16         | 1            | -      | -      |
| 310b  | 85   | 77   | 3.1    | 19      | 23         | 1            | -      | -      |
| 311b  | 78   | 70   | 2.1    | 14      | 16         | 1            | -      | -      |
| 312b  | 76   | 67   | 1.9    | 17      | 23         | 1            | -      | -      |
| 313b  | 71   | 63   | 1.7    | 14      | 15         | 1            | -      | -      |
| 314b  | 80   | 70   | 2.2    | 17      | 21         | 1            | -      | -      |
| 315b  | 77   | 68   | 2.1    | 13      | 15         | 1            | -      | -      |
| 316b  | 93   | 84   | 3.6    | 16      | 23         | 4            | -      | 6      |
| 317b  | 101  | 90   | 4.6    | 11      | 21         | 3            | -      | 6      |
| 318b  | 88   | 79   | 3.3    | 16      | 22         | 4            | 6      | 1      |
| 319b  | 81   | 71   | 2.4    | 19      | 21         | 1            | -      | -      |
| 320b  | 85   | 76   | 2.8    | 20      | 23         | 1            | -      | -      |
| 321b  | 87   | 77   | 3.3    | -       | -          | 6            | 5      | -      |
| 322b  | 80   | 70   | 2.6    | 19      | 21         | 1            | -      | -      |
| 323b  | 73   | 64   | 1.8    | 12      | 13         | 1            | -      | -      |
| 324b  | 79   | 69   | 2.4    | 21      | 24         | 1            | -      | -      |
| 325b  | 76   | 67   | 1.9    | 17      | 20         | 1            | -      | -      |
| 326b  | 84   | 74   | 2.8    | 20      | 21         | 1            | -      | -      |
| 328b  | 78   | 68   | 1.9    | 18      | 19         | 1            | -      | -      |

| Label | TL<br>(cm) | FL<br>(cm) | Weight<br>(kg) | Counted age | Calculated age | Reproductive stage | Egg<br>number | Embryo<br>number |
|-------|------------|------------|----------------|-------------|----------------|--------------------|---------------|------------------|
| 329b  | 87         | 77         | 3.3            | 16          | 21             | <b>1</b>           | -             | -                |
| 330b  | 84         | 75         | 2.8            | 13          | 15             | 5                  | 6             | 5                |
| 331b  | 88         | 78         | 3.5            | 27          | 32             | 4                  | 5             | 7                |
| 332b  | 95         | 85         | 3.9            | 13          | 20             | 2                  | 7             | -                |
| 333b  | 72         | 63         | 1.6            | 12          | 12             | -                  | -             | -                |
| 334b  | 87         | 75         | 2.9            | 15          | 25             | -                  | -             | -                |
| 335b  | 82         | 72         | 2.6            | 18          | 21             | 4                  | -             | 3                |
| 336b  | 85         | 73         | 2.8            | 21          | 23             | 1                  | -             | -                |
| 337b  | 95         | 87         | 4.9            | 17          | 23             | -                  | -             | -                |
| 338b  | 96         | 87         | 4.5            | 12          | 21             | -                  | -             | -                |
| 339b  | 97         | 88         | 4.2            | 18          | 24             | -                  | -             | -                |
| 347b  | 82         | 73         | 2.7            | 17          | 23             | 5                  | 3             | 5                |
| 348b  | 104        | 93         | 5.4            | 17          | 17             | 5                  | 7             | 3                |
| 349b  | 97         | 86         | 4.8            | 14          | 22             | 5                  | 7             | 5                |
| 350b  | 99         | 88         | 5.4            | 10          | 18             | 5                  | ĥ             | 2                |
| 351b  | 78         | 69         | 2.3            | 18          | 20             | •                  |               | 2                |
| 352h  | 87         | 76         | 3.4            | 23          | 21             | 4                  | _             | -<br>A           |
| 353h  | 93         | 82         | 33             | 13          | 20             | 2                  | _             | 6                |
| 360b  | 101        | 88         | 5.1            | 12          | 24             | 5                  | 8             | ă                |
| 361b  | 109        | 97         | 6.9            | 13          | 26             | 5                  | 7             | Ĕ                |
| 362h  | 93         | 83         | 4.2            | 15          | 18             | 5                  | 7             | 4                |
| 363b  | 100        | 90         | 4.6            | 17          | 29             | 4                  | -             | 7                |
| 364b  | 102        | 91         | 5.5            | 17          | 28             | 5                  | 10            | Ŕ                |
| 365h  | 109        | 97         | 7.5            | 12          | 25             | с<br>с             | 11            | 10               |
| 366h  | 91         | 80         | 37             | 21          | 25             | 5                  |               | 10               |

| Appendix 1: continue |
|----------------------|
|----------------------|

| Label | TL   | FL   | Weight        | Counted | Calculated | Reproductive | Egg    | Embryo |
|-------|------|------|---------------|---------|------------|--------------|--------|--------|
|       | (CM) | (cm) | (K <u>G</u> ) | age     | age        | stage        | number | number |
| 367b  | 97   | 86   | 4.3           | 21      | 27         | 4            | 10     | 5      |
| 368b  | 96   | 85   | 4.4           | 12      | 21         | 5            | 7      | 6      |
| 369b  | 97   | 86   | 5.0           | 21      | 28         | 5            | 8      | 7      |

#### Appendix 2: Raw data for male spiny dogfish captured in the Minas Basin from July 7 - October 10, 1996.

| Label       | TL   | FL   | Weight | Counted | Calculated |
|-------------|------|------|--------|---------|------------|
|             | (cm) | (cm) | (kg)   | age     | age        |
| 126b        | 79   | 68   | 1.7    | 15      | 15         |
| 327b        | 83   | 72   | 2.1    | 12      | 15         |
| 340b        | 75   | 66   | 1.7    | 16      | 19         |
| 345b        | 77   | 68   | 1.9    | 14      | -          |
| 346b        | 75   | 66   | 2.1    | •       | -          |
| 354b        | 86   | 76   | 2.6    | 14      | 17         |
| 355b        | 85   | 76   | 2.3    | 10      | 20         |
| 356b        | 75   | 66   | 1.9    | 18      | 21         |
| 357b        | 78   | 68   | 2.0    | 12      | 17         |
| 358b        | 80   | 71   | 2.1    | 15      | 21         |
| <u>359b</u> | 74   | 65   | 2.0    | 16      | 17         |

TL = Total length; FL = Fork length.

# Appendix 3: Raw data for female spiny dogfish captured during the N246 Scotian Shelf DFO survey of July 1996.

| Label | TL<br>(cm) | FL<br>(cm) | Weight<br>(kg) | Counted<br>age | Calculated age | Reproductive<br>stage | Egg<br>number | Embryo<br>number |
|-------|------------|------------|----------------|----------------|----------------|-----------------------|---------------|------------------|
| 1a    | 67         | 57         | 1.135          | 13             | 13             | 1                     |               | -                |
| 2a    | 72         | 63         | 1.345          | 10             | 13             | 1                     | -             | -                |
| 3a    | 77         | 68         | 1.785          | 15             | 19             | 1                     | -             | -                |
| 4a    | 64         | 56         | 1.005          | 11             | -              | 1                     | _             | -                |
| 5a    | 80         | 71         | 2.035          | 17             | 19             | 1                     | -             | -                |
| 6a    | 83         | 72         | 2.250          | 14             | 17             | 1                     | -             | -                |
| 7a    | 67         | 59         | 1.265          | 14             | 14             | 1                     | -             | -                |
| 8a    | 57         | 50         | 0.630          | 4              | 4              | 1                     | -             | _                |
| 9a    | 70         | 60         | 1.295          | 13             | 14             | 1                     | -             | -                |
| 10a   | 82         | 72         | 2.340          | 15             | 18             | 2                     | 4             | -                |
| 21a   | 57         | 50         | 0.895          | 9              | 9              | - 1                   | -             | _                |
| 22a   | 63         | 56         | 1.050          | 12             | 12             | 1                     | _             | _                |
| 23a   | 60         | 53         | 0.905          | 11             | 11             | 1                     | -             | -                |
| 24a   | 68         | 60         | 1.335          | 14             | 14             | 1                     | -             | _                |
| 25a   | 79         | 69         | 2.160          | 15             | 19             | 1                     | -             | -                |
| 26a   | 81         | 71         | 2.650          | 35             | 36             | 1                     | -             | -                |
| 27a   | 72         | 63         | 2.340          | 19             | 21             | i                     | -             | -                |
| 28a   | 63         | 56         | 1.015          | 9              | 9              | i                     | _             | _                |
| 29a   | 53         | 46         | 0.630          | 10             | 10             | 1                     | -             | _                |
| 30a   | 70         | 62         | 1.345          | 16             | 17             | 1                     | _             | _                |
| 51a   | 78         | 68         | 2.202          | 21             | 26             | 2                     | 5             | -                |
| 52a   | 84         | 74         | 2.606          | 14             | 18             | 5                     | 5             | -                |
| 53a   | 78         | 68         | 2.034          | 12             | 15             | 5                     | -             | -+               |
| 54a   | 75         | 67         | 1.714          | 20             | 25             | - 1                   | -             | -                |
| 558   | 70         | 61         | 1 416          | 20             | 20             | 1                     | -             | -                |

TL = Total length; FL = Fork length.

| Label | TL   | FL        | Weight | Counted | Calculated | Reproductive | Egg    | Embryo |
|-------|------|-----------|--------|---------|------------|--------------|--------|--------|
|       | (cm) | (cm)      | (kg)   | age     | age        | stage        | number | number |
| 56a   | 74   | 66        | 1.920  | 15      | 16         | 1            |        | -      |
| 57a   | 67   | 58        | 1.188  | 16      | 16         | 1            | -      | -      |
| 58a   | 83   | 75        | 2.532  | 17      | 22         | 1            | -      | -      |
| 59a   | 81   | 72        | 2.240  | 19      | 25         | 1            | -      | -      |
| 60a   | 58   | 52        | 0.762  | 8       | 8          | 1            | -      | -      |
| 61a   | 77   | 68        | 1.876  | 13      | 18         | -            | -      | -      |
| 62a   | 69   | 61        | 1.430  | 9       | 9          | -            | -      | -      |
| 63a   | 77   | 68        | 1.812  | 11      | 13         | 1            | -      | -      |
| 64a   | 67   | <b>59</b> | 1.354  | 17      | 18         | -            | -      | -      |
| 65a   | 76   | 67        | 2.028  | 17      | 19         | -            | -      | -      |
| 66a   | 73   | 65        | 1.688  | 17      | 17         | -            | -      | -      |
| 75a   | 68   | 60        | 1.365  | 12      | 15         | 1            | -      | -      |
| 76a   | 77   | 67        | 1.885  | 15      | 18         | 1            | -      | -      |
| 77a   | 74   | 67        | 1.760  | 13      | 18         | 1            | -      | -      |
| 78a   | 55   | 48        | 0.720  | 6       | 6          | 1            | -      | -      |
| 79a   | 74   | 65        | 2.140  | 11      | 14         | 1            | -      | -      |
| 88a   | 82   | 72        | 2.355  | 14      | 15         | 1            | -      | -      |
| 89a   | 74   | 65        | 1.620  | 17      | 21         | 1            | -      | -      |
| 90a   | 72   | 63        | 1.640  | 12      | 12         | 1            | -      | -      |
| 94a   | 74   | 63        | 1.415  | 13      | 14         | 1            | -      | -      |
| 97a   | 81   | 73        | 2.290  | 16      | 22         | 1            | -      | _      |
| 98a   | 66   | 58        | 1.215  | 10      | 10         | 1            | -      | _      |
| 99a   | 77   | 68        | 1.885  | 13      | 13         | 1            | -      | -      |
| 111a  | 74   | 64        | 1.870  | 12      | 18         | í            | -      | -      |
| 112a  | 59   | 51        | 0.825  | 8       | 8          | 1            | -      | -      |

| Label | TL   | FL   | Weight | Counted | Calculated | Reproductive | Egg    | Embryo  |
|-------|------|------|--------|---------|------------|--------------|--------|---------|
|       | (cm) | (cm) | (kg)   | age     | age        | stage        | number | number  |
| 113a  | 72   | 63   | 1.450  | 18      | 20         | 1            | -      | -       |
| 114a  | 74   | 65   | 1.745  | 16      | 19         | 1            | -      | -       |
| 115a  | 69   | 61   | 1.395  | 14      | 16         | 1            | -      | -       |
| 116a  | 87   | 76   | 3.155  | 15      | 22         | 5            | 5      | 5       |
| 117a  | 72   | 64   | 1.370  | 20      | 23         | 1            | -      | -       |
| 118a  | 79   | 70   | 2.245  | 11      | 17         | 1            | -      | -       |
| 119a  | 71   | 62   | 1.435  | 20      | 22         | 1            | -      | -       |
| 120a  | 80   | 71   | 2.320  | 16      | 17         | 1            | -      | -       |
| 121a  | 77   | 69   | 1.855  | 17      | 20         | -            | -      | -       |
| 130a  | 69   | 61   | 1.460  | 17      | 19         | 1            | -      | -       |
| 132a  | 68   | 60   | 1.355  | 13      | 13         | 1            | -      | -       |
| 138a  | 66   | 58   | 1.190  | 13      | 13         | 1            | -      | -       |
| 139a  | 74   | 65   | 1.600  | 23      | 26         | 1            | -      | -       |
| 146a  | 69   | 61   | 1.310  | 30      | 31         | 1            | -      | -       |
| 147a  | 78   | 68   | 2.070  | 20      | 24         | 1            | -      | -       |
| 154a  | 70   | 62   | 1.665  | 16      | 21         | 1            | -      | -       |
| 157a  | 57   | 51   | 0.760  | 11      | 11         | 1            | -      | -       |
| 158a  | 65   | 57   | 1.155  | 13      | 13         | 1            | -      | -       |
| 159a  | 67   | 58   | 1.165  | 11      | 1          | 1            | -      | -       |
| 160a  | 61   | 53   | 0.955  | 12      | 12         | 1            | -      | -       |
| 161a  | 74   | 66   | 1.680  | 12      | 14         | 1            | -      | -       |
| 162a  | 65   | 57   | 1.195  | 12      | 12         | -            | -      | -       |
| 1639  | 63   | 55   | 0.990  | 11      | 11         | 1            | -      | -       |
| 164a  | 80   | 71   | 2.210  | 17      | 20         | 1            | -      | -       |
| 165a  | 77   | 68   | 2.175  | 18      | 20         | 1            | -      | <u></u> |

| Label        |        | 167a  | 177a     | 183a     | 184a   |   | 1828  | 186a     | 193a         | 194a     | 195a     | 196a  | 198a       | 199a  | 205a         | 210a       | 212a  | 215a   |
|--------------|--------|-------|----------|----------|--------|---|-------|----------|--------------|----------|----------|-------|------------|-------|--------------|------------|-------|--------|
| ᅻ            | (cm)   | 68    | 63       | 75       | 59     | 3 | 63    | 71       | 67           | 52       | 67       | 64    | 74         | 58    | 51           | 29         | 30    | 29     |
| FL           | (cm)   | 61    | 55       | 67       | л<br>- |   | 55    | 63       | 59           | 46       | 58       | 55    | 64         | 51    | 44           | 26         | 26    | 25     |
| Weight       | (kg)   | 1.300 | 1.160    | 1.855    | 0 870  |   | 0.945 | 1.490    | 1.315        | 0.565    | 1.410    | 1.020 | 1.535      | 0.760 | 0.455        | 0.085      | 0.105 | 0.090  |
| Counted      | age    | 14    |          | 1        | 7 4    | - | 12    | 17       | 17           |          | 10       | 15    | 12         | 10    | 8            | 4          | ധ     | ى<br>س |
| Calculated   | age    | 14    |          | •        | *      | 4 | 12    | 17       | 17           |          | 10       | 15    | 14         | 10    | <b>CB</b> ;  | <b>A</b> : | . دى  | ، دن   |
| Reproductive | stage  | 4     | <b>.</b> | <b>-</b> | • -    | _ |       | <b>_</b> | <b>-\$</b> . | <b>.</b> | <b>.</b> |       | <b>.</b>   |       | <b>-</b> • . | <b>.</b>   | ➡ .   | - ▲    |
| Eaa          | number | _     | 1 1      | •        | •      | r | •     | •        | •            | I        | 1        | ı     | <b>I</b> 1 | •     | •            | L          |       | 1 1    |
| Embryo       | number |       | ı        | ,        | •      | • | •     | I        |              | I 1      | 1 1      | ι ι   | 1          |       | •            | 1          | : 1   | ı      |

Appendix 3: continued

### Appendix 4: Raw data for male spiny dogfish captured during the N246 Scotian Shelf DFO survey of July 1996. TL =

| Label        | TL   | FL   | Weight | Counted | Calculated |
|--------------|------|------|--------|---------|------------|
|              | (cm) | (cm) | (kg)   | age     | age        |
| 11a          | 83   | 73   | 2.010  | 17      | 21         |
| 12a          | 77   | 68   | 1.820  | 13      | 16         |
| 13a          | 67   | 39   | 1.120  | 11      | 11         |
| 1 <b>4</b> a | 67   | 59   | 1.155  | 11      | 11         |
| 15a          | 71   | 62   | 1.360  | 27      | 29         |
| 16a          | 56   | 49   | 0.725  | 8       | 8          |
| 17a          | 68   | 60   | 1.210  | 11      | 11         |
| 18a          | 75   | 67   | 1.605  | 17      | 19         |
| 19a          | 73   | 65   | 1.585  | 5       | 16         |
| 20a          | 65   | 57   | 1.105  | -       | -          |
| 31a          | 77   | 67   | 1.955  | 14      | 17         |
| 32a          | 68   | 61   | 1.355  | 11      | 12         |
| 33a          | 74   | 66   | 1.625  | 16      | 18         |
| 34a          | 55   | 49   | 0.730  | 9       | 9          |
| 35a          | 63   | 55   | 1.105  | 5       | 9          |
| 36a          | 69   | 61   | 1.340  | 15      | 15         |
| 37a          | 66   | 58   | 1.220  | 11      | 11         |
| 38a          | 67   | 58   | 1.180  | 12      | 12         |
| 39a          | 78   | 68   | 1.800  | 11      | 17         |
| 40a          | 62   | 55   | 1.005  | 7       | 13         |
| 41a          | 85   | 74   | 2.416  | 7       | 14         |
| 42a          | 72   | 64   | 1.582  | -       | -          |
| <b>4</b> 3a  | 62   | 55   | 0.984  | 13      | 13         |
| 44a          | 79   | 71   | 2.176  | 13      | 25         |
| 45a          | 76   | 66   | 1.730  | 11      | 15         |

Total length; FL = Fork length.

| Label        | TL   | FL   | Weight | Counted | Calculated |
|--------------|------|------|--------|---------|------------|
|              | (cm) | (cm) | (kg)   | age     | age        |
| 46a          | 69   | 56   | 1.262  | 10      | 16         |
| 47a          | 61   | 54   | 0.942  | 9       | 9          |
| 48a          | 86   | 77   | 2.416  | 8       | 19         |
| 49a          | 65   | 57   | 1.110  | 12      | 12         |
| 50a          | 66   | 58   | 1.248  | 13      | 13         |
| 67a          | 79   | 69   | 2.176  | 8       | 17         |
| 68a          | 77   | 68   | 1.730  | 7       | 13         |
| 69a          | 67   | 59   | 1.274  | 12      | 12         |
| 70a          | 72   | 64   | 1.394  | 18      | -          |
| 71a          | 74   | 65   | 1.662  | 10      | 12         |
| 72a          | 72   | 64   | 1.396  | 14      | 18         |
| 73a          | 76   | 57   | 1.704  | 15      | 19         |
| 7 <b>4</b> a | 69   | 60   | 1.322  | 11      | 11         |
| 80a          | 69   | 61   | 1.480  | 15      | 17         |
| 81a          | 64   | 56   | 0.995  | 13      | 13         |
| 82a          | 74   | 64   | 1.735  | 11      | 21         |
| 83a          | 73   | 64   | 1.565  | 8       | 8          |
| 84a          | 65   | 58   | 1.115  | 9       | 12         |
| 85a          | 84   | 75   | 2.255  | -       | -          |
| 86a          | 70   | 62   | 1.555  | 15      | 15         |
| 87a          | 71   | 62   | 1.835  | 10      | 13         |
| 91a          | 77   | 67   | 1.715  | 17      | 22         |
| 92a          | 71   | 62   | 1.350  | 6       | 10         |
| 93a          | 74   | 65   | 1.660  | 15      | 20         |
| 95a          | 83   | 74   | 2.330  | 11      | 16         |

| 1368  |                | 1359  | 134a   | 133a  | 131a  | 129a  | 128a  | 127a  | 126a  | 125a  | 124a  | 123a     | 122a  | 110a  | 109a     | 108a  | 107a  | 106a  | 105a  | 104a  | 103a  | 102a  | 101a  | 100a  | 96a   |      | Label      | ***** |
|-------|----------------|-------|--------|-------|-------|-------|-------|-------|-------|-------|-------|----------|-------|-------|----------|-------|-------|-------|-------|-------|-------|-------|-------|-------|-------|------|------------|-------|
| 12    | 1 <del>-</del> | 74    | 75     | 78    | 74    | 84    | 82    | 75    | 77    | 69    | 71    | 77       | 64    | 68    | 69       | 70    | 69    | 75    | 65    | 67    | 77    | 79    | 68    | 71    | 75    | (cm) | ר<br>ב     |       |
| 63    | CO             |       | 66     | 68    | 65    | 73    | 73    | 66    | 68    | 62    | 63    | 68       | 57    | 69    | 61       | 62    | 61    | 68    | 57    | 59    | 68    | 70    | 59    | 62    | 66    | (cm) | FL         |       |
| 1.575 | 1.860          | 1.010 | 1 540  | 1.845 | 1.550 | 2.515 | 2.015 | 1.545 | 1.595 | 1.300 | 1.405 | 1.910    | 0.990 | 1.780 | 1.140    | 1.460 | 1.250 | 1.865 | 1.245 | 1.395 | 1.890 | 1.875 | 1.285 | 1.530 | 1.640 | (kg) | Weight     |       |
| 19    | 20             | 32    | 19     | 23    | 16    | 20    | 10    | 15    | 11    | 16    | 17    | <u>-</u> | 14    | 12    | 11       | 16    | 18    | ı     | 13    | 16    | 18    | 15    | 17    | 10    | 7     | age  | Counted    |       |
| 21    | 23             | 20    | 5<br>ī | 27    | 16    | 27    | 23    | 16    | 14    | 16    | 17    | 15       | 14    | 100   | <b>-</b> | 10    | 17    | •     |       | 17    | 21    | 18    | 17    | 14    | 10    | age  | Calculated |       |

| Label | TL   | FL   | Weight | Counted | Calculated |
|-------|------|------|--------|---------|------------|
|       | (cm) | (cm) | (kg)   | age     | age        |
| 137a  | 70   | 61   | 1.345  | 15      | 15         |
| 140a  | 78   | 69   | 1.780  | 13      | 17         |
| 141a  | 71   | 62   | 1.455  | 13      | 19         |
| 142a  | 80   | 71   | 1.870  | 8       | -          |
| 143a  | 77   | 69   | 1.865  | 11      | 15         |
| 144a  | 81   | 72   | 2.210  | 10      | 19         |
| 145a  | 66   | 58   | 1.195  | 17      | 17         |
| 148a  | 74   | 66   | 1.755  | 23      | 23         |
| 149a  | 63   | 56   | 1.005  | 19      | 19         |
| 150a  | 69   | 61   | 1.335  | 15      | 16         |
| 151a  | 73   | 65   | 1.580  | 19      | 26         |
| 152a  | 68   | 60   | 1.150  | 17      | 17         |
| 153a  | 79   | 71   | 2.095  | 21      | 25         |
| 155a  | 64   | 57   | 1.025  | 17      | 17         |
| 156a  | 74   | 65   | 1.470  | 14      | 15         |
| 162a  | 65   | 57   | 1.195  | 12      | 12         |
| 166a  | 79   | 70   | 2.000  | 12      | 15         |
| 168a  | 79   | 70   | 2.270  | 13      | 18         |
| 169a  | 66   | 59   | 1.135  | 15      | 15         |
| 170a  | 77   | 68   | 1.730  | 16      | 16         |
| 171a  | 71   | 62   | 1.590  | 16      | 17         |
| 172a  | 71   | 63   | 1.630  | 15      | 16         |
| 173a  | 72   | 64   | 1.390  | 10      | 10         |
| 174a  | 69   | 61   | 1.310  | 12      | 13         |
| 175a  | 66   | 59   | 1.130  | 9       | 9          |

| Label         | TL   | FL   | Weight | Counted | Calculated |
|---------------|------|------|--------|---------|------------|
|               | (cm) | (cm) | (kg)   | age     | age        |
| 176a          | 61   | 54   | 0.900  | 12      | 12         |
| 178a          | 63   | 55   | 0.935  | 10      | 10         |
| 179a          | 73   | 63   | 1.510  | 13      | 14         |
| 180a          | 71   | 63   | 1.235  | 16      | 17         |
| 181a          | 73   | 65   | 1.525  | 11      | 11         |
| <b>182</b> a  | 71   | 63   | 1.450  | 14      | 15         |
| 187a          | 75   | 66   | 1.590  | 13      | 13         |
| 188a          | 76   | 67   | 1.620  | -       | -          |
| 189a          | 53   | 46   | 0.645  | 6       | 6          |
| 190a          | 57   | 50   | 0.775  | 8       | 8          |
| 191a          | 56   | 49   | 0.600  | 6       | 6          |
| 1 <b>92</b> a | 71   | 63   | 1.335  | 12      | 14         |
| 197a          | 69   | 60   | 1.255  | 17      | 20         |
| 200a          | 74   | 64   | 1.405  | 13      | 14         |
| 201a          | 77   | 68   | 1.875  | 10      | 13         |
| 202a          | 73   | 65   | 1.455  | 13      | 13         |
| 203a          | 69   | 61   | 1.070  | 11      | 11         |
| 204a          | 64   | 57   | 0.925  | 10      | 10         |
| 206a          | 77   | 68   | 1.700  | 6       | 9          |
| 207a          | 60   | 54   | 0.925  | 10      | 10         |
| 208a          | 76   | 67   | 2.160  | -       | -          |
| 209a          | 69   | 62   | 1.225  | 9       | 11         |
| 211a          | 25   | 29   | 0.075  | 2       | 2          |
| 213a          | 25   | 28   | 0.075  | 3       | 3          |
| 214a          | 37   | 41   | 0.280  | 4       | 4          |

#### Appendix 5: Raw data for female spiny dogfish captured during the N249 Gulf of St. Lawrence DFO survey of

| Label | TL   | FL   | Weight | Counted | Calculated | Reproductive | Egg    | Embryo |
|-------|------|------|--------|---------|------------|--------------|--------|--------|
|       | (cm) | (cm) | (kg)   | age     | age        | stage        | number | number |
| 2d    | 67   | 60   | 1.285  | 14      | 14         | 1            |        | -      |
| 3d    | 86   | 77   | 2.756  | 20      | 26         | 1            | -      | -      |
| 4d    | 76   | 66   | 1.886  | 15      | 22         | 1            | -      | -      |
| 5d    | 70   | 62   | 1.188  | 13      | 12         | 1            | -      | -      |
| 8d    | 87   | 77   | 2.306  | 18      | 22         | 1            | **     | -      |
| 9d    | 87   | 77   | 2.920  | 13      | 13         | 5            | 3      | 5      |
| 20d   | 85   | 76   | 2.870  | 12      | 18         | 6            | 4      | -      |
| 21d   | 90   | 79   | 3.26   | 17      | 18         | 3            | -      | 5      |
| 22d   | 85   | 76   | 2.88   | 17      | 21         | 1            | -      | -      |
| 23d   | 101  | 89   | 5.12   | 11      | 15         | 3            | -      | 7      |
| 24d   | 64   | 57   | 1.18   | 12      | 14         | 1            | -      | -      |
| 25d   | 73   | 64   | 1.64   | 11      | 14         | 1            | -      | -      |
| 26d   | 80   | 70   | 2.35   | 14      | 17         | 1            | -      | -      |
| 27d   | 66   | 56   | 1.31   | 9       | 9          | 1            | -      | -      |
| 28d   | 88   | 78   | 3.41   | 15      | 20         | 3            | -      | 4      |
| 29d   | 93   | 82   | 3.98   | 16      | 18         | 3            | -      | 6      |
| 30d   | 97   | 86   | 3.86   | 12      | 17         | 3            | -      | 5      |
| 31d   | 98   | 86   | 4.27   | -       | -          | 1            | -      | -      |
| 32d   | 95   | 84   | 4.18   | 12      | 25         | 3            | -      | 7      |
| 33d   | 76   | 67   | 1.59   | 18      | 20         | 1            | -      | -      |
| 35d   | 92   | 82   | 3.67   | 15      | 21         | 3            | -      | 3      |
| 36d   | 93   | 81   | 3.12   | 12      | 18         | 1            | -      | -      |
| 37d   | 89   | 78   | 2.50   | 14      | 20         | 1            | -      | -      |
| 38d   | 79   | 69   | 1.86   | 11      | 16         | 1            | -      | -      |
| 39d   | 82   | 73   | 2.11   | 17      | 19         | 1            | -      | -      |

September 1996. TL = Total length; FL = Fork length.

| Label | TL   | FL          | Weight      | Counted | Calculated | Reproductive | Egg    | Embryo |
|-------|------|-------------|-------------|---------|------------|--------------|--------|--------|
|       | (cm) | <u>(cm)</u> | <u>(kg)</u> | age     | age        | stage        | number | number |
| 40d   | 81   | 71          | 2.23        | 13      | 19         | 3            | -      | 3      |
| 41d   | 79   | 69          | 2.02        | 12      | 17         | 1            | -      | -      |
| 42d   | 81   | 71          | 2.24        | 14      | 17         | 1            | -      | -      |
| 43d   | 81   | 71          | 2.11        | 11      | 13         | 1            | -      | -      |
| 44d   | 82   | 72          | 2.32        | 8       | 15         | 1            | -      | -      |
| 51d   | 93   | 82          | 3.67        | -       | -          | 4            | -      | 4      |
| 52d   | 67   | 59          | 1.16        | 14      | 14         | 1            | -      | -      |
| 53d   | 88   | 78          | 2.89        | 16      | 21         | 1            | -      | -      |
| 54d   | 84   | 74          | 2.15        | 15      | 16         | 1            | -      | -      |
| 55d   | 78   | 68          | 1.71        | 12      | 13         | 1            | -      | -      |
| 56d   | 78   | 68          | 1.70        | 13      | 13         | 1            | -      | -      |
| 63d   | 88   | 78          | 2.92        | 21      | 23         | 1            | -      | -      |
| 64d   | 85   | 75          | 3.23        | 12      | 17         | 3            | -      | 3      |
| 65d   | 96   | 85          | 3.70        | 18      | 23         | 1            | -      | -      |
| 66d   | 87   | 77          | 3.38        | 17      | 22         | 5            | 6      | 3      |
| 69d   | 84   | 75          | 2.39        | 22      | 27         | -            | -      | -      |
| 70d   | 86   | 76          | 2.89        | 21      | 23         | 3            | -      | 3      |
| 71d   | 75   | 67          | 1.97        | 12      | 18         | 1            | -      | -      |
| 72d   | 91   | 81          | 3.41        | 16      | 21         | 3            | -      | 3      |
| 73d   | 89   | 79          | 3.00        | 17      | 22         | 5            | 4      | 3      |
| 74d   | 91   | 80          | 3.38        | 18      | 22         | 6            | 1      | -      |
| 75d   | 88   | 78          | 2.65        | 17      | 24         | 1            | -      | -      |
| 76d   | 81   | 72          | 2.32        | 18      | 19         | 1            | -      | -      |
| 77d   | 82   | 72          | 2.56        | 18      | 20         | 1            | -      | -      |
| 80d   | 86   | 76          | 3.06        | -       | -          | 1            | -      | -      |

| Label | TL   | FL   | Weight | eight Counted Calculated Repr |     | Reproductive | Egg    | Embryo |
|-------|------|------|--------|-------------------------------|-----|--------------|--------|--------|
|       | (cm) | (cm) | (kg)   | age                           | age | stage        | number | number |
| 82d   | 78   | 68   | 1.70   | 16                            | 18  | 1            | -      | -      |
| 83d   | 82   | 71   | 2.41   | 10                            | 17  | 6            | 6      | -      |
| 84d   | 77   | 67   | 2.02   | 12                            | 13  | 1            | -      | -      |
| 86d   | 65   | 56   | 1.08   | 10                            | 10  | 1            | -      | -      |
| 93d   | 77   | 67   | 1.40   | 12                            | 21  | 1            | -      | -      |
| 94d   | 78   | 68   | 1.90   | 20                            | 22  | 1            | -      | -      |
| 95d   | 73   | 63   | 1.70   | 17                            | 19  | 1            | -      | -      |
| 96d   | 76   | 66   | 1.50   | 17                            | 18  | 1            | _      | _      |
| 97d   | 80   | 70   | 2.20   | 16                            | 19  | 1            | -      | -      |
| 98d   | 75   | 65   | 1.40   | 15                            | 16  | 1            | -      | -      |
| 99d   | 79   | 68   | 1.80   | 21                            | 22  | 1            | -      | -      |
| 109d  | 78   | 68   | 2.17   | 15                            | 20  | 1            | -      | -      |
| 110d  | 78   | 68   | 2.17   | 15                            | 15  | 1            | -      | -      |
| 111d  | 64   | 56   | 1.16   | 15                            | 15  | 1            | -      | -      |
| 112d  | 64   | 55   | 0.90   | 19                            | 22  | 1            | -      | -      |
| 113d  | 77   | 68   | 1.80   | -                             | -   | 1            | -      | -      |
| 114d  | 75   | 65   | 1.80   | 17                            | 19  | 1            | -      | -      |

### Appendix 6: Raw data for male spiny dogfish captured during the N249 Gulf of St. Lawrence DFO survey of September

| Label | TL   | FL   | Weight | Counted | Calculated |
|-------|------|------|--------|---------|------------|
|       | (cm) | (cm) | (kg)   | age     | age        |
| 1d    | 75   | 64   | 1.596  | 21      | 23         |
| 6d    | 87   | 77   | 2.506  | 5       | 16         |
| 7d    | 72   | 63   | 1.516  | 21      | 22         |
| 10d   | 70   | 60   | 1.540  | 11      | 12         |
| 11d   | 74   | 65   | 1.64   | 13      | 17         |
| 12d   | 79   | 69   | 1.96   | 12      | 14         |
| 13d   | 81   | 70   | 2.05   | 13      | 16         |
| 14d   | 77   | 68   | 1.97   | 15      | 16         |
| 15d   | 79   | 69   | 2.39   | 9       | 10         |
| 16d   | 75   | 65   | 1.88   | 3       | 12         |
| 17d   | 69   | 61   | 1.37   | 11      | 16         |
| 18d   | 76   | 67   | 1.77   | 7       | 15         |
| 19d   | 81   | 71   | 1.83   | 5       | 23         |
| 34d   | 84   | 74   | 1.98   | 17      | 21         |
| 45d   | 75   | 66   | 1.54   | 11      | 13         |
| 46d   | 76   | 67   | 1.62   | -       | -          |
| 47d   | 78   | 69   | 2.12   | 10      | 13         |
| 48d   | 78   | 68   | 1.87   | 13      | 15         |
| 49d   | 84   | 74   | 2.15   | 10      | 17         |
| 50d   | 73   | 63   | 1.37   | 10      | 11         |
| 57d   | 79   | 69   | 1.70   | 13      | 16         |
| 58d   | 80   | 70   | 1.80   | 12      | 18         |
| 59d   | 79   | 69   | 1.67   | 11      | 14         |
| 60d   | 71   | 62   | 1.39   | 17      | 17         |
| 61d   | 71   | 63   | 1.07   | 18      | 18         |

1996. TL = Total length; FL = Fork length.

152

•

| Label | TL   | FL   | Weight | Counted | Calculated |
|-------|------|------|--------|---------|------------|
|       | (cm) | (cm) | (kg)   | age     | age        |
| 62d   | 70   | 61   | 1.17   | 12      | 12         |
| 67d   | 79   | 71   | 2.09   | 11      | 13         |
| 68d   | 78   | 69   | 1.98   | 7       | 14         |
| 79d   | 77   | 67   | 1.88   | 16      | 21         |
| 81d   | 68   | 59   | 1.10   | 10      | 10         |
| 85d   | 69   | 61   | 1.30   | 9       | 9          |
| 87d   | 78   | 68   | 1.88   | 10      | 14         |
| 88d   | 74   | 64   | 1.43   | 12      | 14         |
| 89d   | 72   | 64   | 1.37   | 9       | 13         |
| 90d   | 65   | 57   | 0.98   | 13      | 13         |
| 91d   | 77   | 68   | 1.83   | 16      | 19         |
| 92d   | 73   | 63   | 1.40   | 13      | 22         |
| 100d  | 77   | 67   | 1.6    | 15      | 17         |
| 101d  | 75   | 65   | 1.5    | 17      | 21         |
| 102d  | 74   | 64   | 1.5    | 17      | 17         |
| 103d  | 72   | 62   | 1.5    | 13      | 14         |
| 104d  | 75   | 66   | 1.7    | -       | -          |
| 105d  | 75   | 65   | 1.5    | 13      | 16         |
| 106d  | 82   | 72   | 2.3    | -       | -          |
| 107d  | 76   | 66   | 1.7    | -       | -          |
| 108d  | 71   | 62   | 1.4    | 23      | 24         |







IMAGE EVALUATION TEST TARGET (QA-3)







C 1993, Applied Image, Inc., All Rights Reserved

